



nKPI Technical Documentation

June 2026

MedicalDirector Insights 1.9.5

**Specifications for National Key
Performance Indicators and Online
Services Reporting 20.0**

Note: This information relates to V20.0 of the *Specifications for National Key Performance Indicators and Online Services Reporting* (Solving Health, Specifications for National Key Performance Indicators and Online Services Reporting V20.0. Sydney: Solving Health, February 2026). For more information, see <https://www.solvinghealth.au/specifications>.

DOCUMENT CONTROL

Version History

Version	Date	Author	Relevant MD Insights release	Changes
1.0	June, 2024	MedicalDirector	MD Insights 1.9.0	Initial draft
1.1	December, 2024	MedicalDirector	MD Insights 1.9.1	Minor updates to text for PI26
1.2	June, 2025	MedicalDirector	MD Insights 1.9.2	- PI05 and PI06: Added new LOINC codes and local codes - PI18 and PI19: Included a free-text search for pathology test names - PI25: Updated LOINC codes - PI26: Added diagnosis codes
1.2	December, 2025	MedicalDirector	MD Insights 1.9.3	- Addition of new local pathology codes for HbA1c, ACR and eGFR - PI07: Inclusion of new MBS item numbers - PI14: Inclusion of additional vaccines - PI18 and PI19 - Review ACR and eGFR LOINC codes

				- PI19: Implement updated uACR thresholds - PI20 and PI21: Added diagnosis codes - PI23 and PI24: Added diagnosis codes - PI25: Addition of free text terms and local pathology codes
1.3	June, 2026	MedicalDirector	MD Insights 1.9.5	- Visit type: Telephone visit type counted as a visit - PI11: Addition of business rule for birthweight - PI13: Addition of business rule for birthweight - PI14: Inclusion of additional vaccines

Contents

DOCUMENT CONTROL	2
INTRODUCTION	5
NATIONAL KEY PERFORMANCE INDICATORS	8
PI01 – BIRTHWEIGHT RECORDED	8
PI02 – BIRTHWEIGHT RESULT	10
PI03 – MBS HEALTH ASSESSMENT	13
PI05 – HbA1c RECORDED (TYPE 2 DIABETES PATIENTS)	15
PI06 – HbA1c RESULTS (TYPE 2 DIABETES PATIENTS)	20
PI07 – CHRONIC DISEASE MANAGEMENT PLAN PREPARED	24
PI09 – SMOKING STATUS RECORDED	27
PI10 – SMOKING STATUS RESULT	29
PI11 – SMOKING DURING PREGNANCY	31
PI12 – BODY MASS INDEX	36
PI13 – FIRST ANTENATAL CARE VISIT	40
PI14 – INFLUENZA IMMUNISATION	43
PI16 – ALCOHOL CONSUMPTION RECORDED	46
PI18 – KIDNEY FUNCTION TEST RECORDED (TYPE 2 DIABETES OR CVD PATIENTS)	48
PI19 – KIDNEY FUNCTION TEST RESULT (TYPE 2 DIABETES OR CVD PATIENTS)	53
PI20 – CARDIOVASCULAR DISEASE (CVD) RISK ASSESSMENT	57
PI21 – CARDIOVASCULAR DISEASE (CVD) RISK ASSESSMENT RESULTS	63
PI22 – CERVICAL SCREENING RECORDED	67
PI23 – BLOOD PRESSURE RECORDED (TYPE 2 DIABETES PATIENTS)	70
PI24 – BLOOD PRESSURE LESS THAN OR EQUAL TO 140/90 MMHG (TYPE 2 DIABETES PATIENTS)	72
PI25 – STI RESULT RECORDED	74
PI26 – EAR HEALTH CHECK	75
APPENDIX A- DOCLE CODES	77
APPENDIX B- LOINC CODES	84
APPENDIX C- FREE TEXT SEARCH TERMS FOR PI25	87
APPENDIX D- CREATING AND PROCESSING A CLAIM	89
EXAMPLE - PROCESSING A PATIENT CLAIM	90
APPENDIX E- RETIRED AND SUSPENDED INDICATORS	93
PI04 – FULLY IMMUNISED CHILDREN (SUSPENDED)	93
PI08 – TEAM CARE ARRANGEMENT (MBS ITEM 723) (RETIRED)	97
PI15 – INFLUENZA IMMUNISATION (TYPE 2 DIABETES OR COPD PATIENTS) (RETIRED)	99
PI17 – AUDIT-C RESULT RECORDED (RETIRED)	101

Introduction


This purpose of this document is to guide you through the process of accurately recording data points during a clinical consult to ensure effective nKPI reports can be created.

Frequent Definitions

Aboriginal Patient Definition

A client who is recorded in the clinical software identifying as Aboriginal or Torres Strait Islander.

Patient Details



Pt. Details | Allergies/Adverse Reactions/Warnings | Family/Social Hx | Notes | Smoking | Alcohol | Personal Details

Title: ☐ Single Name ☐ Head of Family ☐

First Name:

Middle Name:

Surname:

Known as:

Date of Birth: Sex at Birth: ☐ Transgender

Gender Identity:

Pronouns:

ATSI:

- Not Stated
- Aboriginal
- Torres Strait Islander
- Aboriginal and Torres Strait Islander
- Neither Aboriginal nor Torres Strait Islander

Ethnicity:

Contact Details:

Residential:

Address:

City/Suburb: Postcode:

Default phone number to be displayed: ☐ Home ☐ Work ☐ Mobile

Phone:

E-mail:

Preferred Contact: ☐ Do not send SMS

Medicare No:

Medicare Expiry:

Pension No:

DVA No: ☒ Validate

Safety Net No:

Record No:

IHI No:

IHI Record Status:

IHI No Status:

Regular Practitioner:

Access Restriction: ☒ No Restriction ☐ Users with Restricted Access Permission Only ☐ Regular Practitioner Only

☐ This patient is registered with this Practice for MyMedicare.

☐ This patient has withdrawn consent to upload clinical documents to MyHealthRecord (except for prescription records and investigation requests).

☐ This patient has withdrawn consent to upload de-identified clinical data to the MedicalDirector PIP QI data set.

☐ This patient has withdrawn consent to upload de-identified clinical data to the MedicalDirector research data set.

☐ Update address for all family members

☐ Auto-capitalise names

Indigenous status is defined by whether a person identifies as being of Aboriginal or Torres Strait Islander origin.

Permissible values are:

- Aboriginal but not Torres Strait Islander origin
- Torres Strait Islander but not Aboriginal origin
- Both Aboriginal and Torres Strait Islander origin
- Neither Aboriginal nor Torres Strait Islander origin
- Not stated/inadequately described

Note: Categories 1, 2 and 3 are classified as Indigenous.

Category 4 is classified as Non-Indigenous.


The nKPI report includes only Indigenous clients for all the indicators.

Regular Patient Definition

A client who has 3 or more visits within the previous 2 years recorded in the clinical software. Note: This does not include Deceased and Inactive patients.

Visit Type	
Counted as a visit	Not counted as a visit
Home visit consultation	Email
Hospital consultation	Practice admin
Locum service	SMS
Nursing home consultation	
Others	
Surgery consultation	
Telehealth (clinical)	
Telephone	

A Visit can be added as shown below:



Consultation date: 17/06/2024

Previous visits: ALL

Date	Recorded by:	Visit type	Reason for contact	Start	Duration	Medicare item	Ri
06/08/2023...		Surgery Consultation	Cough	09:30:00			
06/09/2023...		Surgery Consultation	Cold - Common	11:30:00			
17/06/2024	Dr A Practitioner	Surgery Consultation		15:04:18			

The following steps are to record a patient as inactive or deceased:

- 1) Search the patient from the list.

Select Patient From List

Enter patient surname, chart number, phone number or D.O.B. (dd/mm/yyyy)

TEST

Include

☐ Inactive patients

☐ Deceased

☐ Next of Kin and Emergency Contact

Search on

☒ Chart number

☐ Phone number

Name	Age	Gender	Chart Number	Address	Phone Number
Test 1121w, Test 1121w	34yrs ...	M		Aa. Alexandria Mc 2004	
Test_1661645_1, Test_1661645_1	68yrs ...	M		Aa. Abbotsham 7315	
Test1687, Test1687	34yrs ...	M		Aa. Melbourne 3000	
Test1692, Test1692	34yrs ...	M		Aa. Melbourne 3000	

Open

Edit

Delete

Add

New

Merge

Status: Active

OK Add New Delete Edit Merge Close

- 2) Right click on the patient.

- 3) Click the required option.

Select Patient From List

Enter patient surname, chart number, phone number or D.O.B. (dd/mm/yyyy)

TEST

Include

☐ Inactive patients

☐ Deceased

☐ Next of Kin and Emergency Contact

Search on

☒ Chart number

☐ Phone number

Name	Age	Gender	Chart Number	Address	Phone Number
Test 1121w, Test 1121w	34yrs ...	M		Aa. Alexandria Mc 2004	
Test_1661645_1, Test_1661645_1	68yrs ...	M		Aa. Abbotsham 7315	
Test1687, Test1687	34yrs ...	M		Aa. Melbourne 3000	
Test1692, Test1692	34yrs ...	M		Aa. Melbourne 3000	

Delete patient

Do you want to:

Mark as inactive

Mark as deceased

Delete permanently

Cancel

Status: Active

OK Add New Delete Edit Merge Close

NATIONAL KEY PERFORMANCE INDICATORS

PI01 – Birthweight Recorded

Description

Proportion of Indigenous babies born within the previous 12 months who attended the organisation more than once whose birthweight has been recorded at your primary health care service.

- The baby is considered Indigenous if one or both parents identify as Indigenous.
- This patient does not need to be defined as a 'Regular' patient.
- If a patient record exists, it will be considered as a live birth.
- Does not include deceased patient records.

Business Rules

- Only live births are included.
- Results arising from measurements conducted outside of the service, that are known by the service, are included in the calculation of this indicator.
- Where a baby does not have a separate client record, and the mother's record is used as a source of birth details, the baby is considered Indigenous if the mother is Indigenous.

Numerator

- Number of Indigenous babies born within the previous 12 months who have had more than one visit and whose birthweight has been recorded at the primary health care service.

Denominator

- Total number of Indigenous babies born within the previous 12 months who have had more than one visit and had a medical record at the primary health care service.

Records Excluded


- Stillborn babies.
- Babies without a medical record of their own at your organisation, even if their information is recorded in their mother's record.

Notes


- The birthweight is to be taken from the baby's client record where available.
- The date of the weight recorded must match the babies' date of birth. (The date of the recorded weight can be modified in the top left of the window as indicated below in the example).

Example – Enter A Birth Weight for A Patient

1. Open a patient's record.
2. Select on **Tools > Tool Box > Weight**.



3. Paediatric percentile charts will be displayed as below.



The screenshot shows the 'Tool Box' window titled 'Paediatric Percentile Charts'. The patient information at the top includes Date: 14/03/2019, Time: 10:01:39 AM, Gender: Female, Age: 5, Height: 107, and Patient ID: 35. The 'Paediatric' tab is selected, and the 'Data' button is highlighted. The 'Current Measurements' section has input fields for Height (cm), Weight (Kg), and BMI. A table with columns Date, Time, Height/Length, Weight, BMI, and Head circ. is displayed. The 'Clear' button is at the bottom left, and 'Print', 'Reference', 'Edit', 'View', 'Save', and 'Close' buttons are at the bottom right.

4. Enter a value for weight.
 - Note that the date/time fields will autofill to **today's** date and time. Modify this to the patient's date of birth.
5. Click **Save**.

PI02 – Birthweight Result

Description

Proportion of Indigenous babies born within the previous 12 months, who attended the organisation more than once whose birthweight results were categorised as one of the following:

Category	Birth Weights
Low	Less than 2,500 grams
Normal	Between 2,500 grams to less than 4,500 grams
High	Equal to or greater than 4,500 grams

Business Rules

- This patient does not need to be recorded as 'Regular'.
- Only live births are included.
- Results arising from measurements conducted outside of the service, that are known by the service, are included in the calculation of this indicator.
- Where a baby does not have a separate client record, and the mother's record is used as a source of birth details, the baby is considered Indigenous if the mother is Indigenous.

Some special rules apply to birthweights below and above certain thresholds, i.e. values of:

- ≤ 9 are treated as kilograms
- < 400 are treated as kilograms but capped to 9,998 grams
- 400 to 9998 are treated as grams
- $> 9,998$ are treated as grams but capped to 9,998

The above rules apply because MedicalDirector Clinical records weight in kilograms. However, to account for the possibility of data in grams making its way into the database (e.g. via a data import from another software provider), these rules have been implemented to attempt to manage this.

Calculation 1: Babies Born with A Low Birthweight

Numerator

The number of babies born within the previous 12 months, who have had more than one visit and whose birthweight results were categorised as low.

Denominator

Number of indigenous babies with a birthweight recorded and meet the below criteria:

- Only live births as defined in [PI01](#).
- The number of babies born within the previous 12 months, who have had more than one visit and whose birthweight has been recorded. Include all babies with a file, regardless of whether they are regular patients.

Calculation 2: Babies Born with A Normal Birthweight

Numerator

The number of babies born within the previous 12 months, who have had more than one visit and whose birthweight results were categorised as normal.

Denominator

Number of indigenous babies with a birthweight recorded and meet the below criteria:

- Only live births as defined in [PI01](#).
- The number of babies born within the previous 12 months, who have had more than one visit and whose birthweight has been recorded. Include all babies with a file, regardless of whether they are regular patients.

Calculation 3: Babies Born with A High Birthweight

Numerator

The number of babies born within the previous 12 months, who have had more than one visit and whose birthweight results were categorised as high.

Denominator

Number of indigenous babies with a birthweight recorded and meet the below criteria:

- Only live births as defined in [PI01](#).
- The number of babies born within the previous 12 months, who have had more than one visit and whose birthweight has been recorded. Include all babies with a file, regardless of whether they are regular patients.

Records Excluded

- Multiple births (including twins). Babies born as part of multiple births are more likely to have a lower birthweight.
- Babies with unknown birthweight.
- Babies who were stillborn.
- Babies born before 20 weeks gestation.
- Babies with a birthweight recorded of less than 400 grams.
- Babies without a medical record of their own, even if their information is recorded in their mother's record.

Notes

- The birthweight is to be taken from the baby's client record where available.
- Where a baby does not have a separate client record, the mother's record may be used as a source of birth details.

Example - Enter A Birth Weight for A Patient

The birthweight for a patient can be entered following the method outlined in [PI01](#).

Check

- Total number of Indigenous babies born with birthweight recorded = low birthweight + normal birthweight + high birthweight babies.

The Denominator is the same for 1, 2 and 3 above.

The Denominator is equal to or less than the Numerator in [PI01](#) (that is, you cannot have more birthweight results than number of babies with birthweights recorded).

PI03 – MBS Health Assessment

Description

Proportion of Indigenous regular clients who have a current completed Indigenous health assessment as indicated by an MBS-rebate Indigenous health assessment:

- In-person MBS-rebate items - MBS items: 715 or 228.
- Telehealth MBS-rebate items - MBS items: 92004 or 92011.

For ages 0–14: an Indigenous health assessment is counted if it was completed within the previous 12 months.
For ages 15+: an Indigenous health assessment is counted if it was completed within the previous 24 months.

Business Rules

- Patient must have had at least one MBS claim where the 'date lodged' falls in the previous 12 or 24 months.
- The claim is counted at the point of claim submission.

Calculation 1: Indigenous regular clients for whom a Medicare Benefits Schedule Health Assessment for Aboriginal and Torres Strait Islander People (MBS Items 715, 228) was claimed

Numerator

Number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]** who have a current completed in person MBS-rebate Indigenous health assessment (MBS items: 715 or 228).

Include in the Numerator **only** those patients whose MBS Health Assessment for Aboriginal and Torres Strait Islander People (MBS 715 or 228) was claimed by **your organisation**. See DOH (2018) for more information.

Denominator

Total number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]**.

Calculation 2: Indigenous regular clients for whom a Medicare Benefits Schedule Health Assessment for Aboriginal and Torres Strait Islander People (MBS Items 92004, 92011) was claimed

Numerator

Number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]** who have a current completed in-person MBS-rebate Indigenous health assessment (MBS items: 92004, 92011)

Include in the Numerator only those patients whose MBS Health Assessment for Aboriginal and Torres Strait Islander People (MBS 92004, 92011) was claimed by your organisation. See DOH (2018) for more information.

Denominator

Total number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]**

Reporting by Age Group

	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Notes

If your organisation does not claim this MBS item but provides an equivalent level of care, such as a comprehensive health Check that cannot be claimed through the MBS.

For information on recording the visit and transmitting the claim to Medicare please refer to the following help files:

- [Recording a visit.](#)
- [Transmitting bulk claims to Medicare.](#)

Check

For each age and sex at birth group, you cannot have more patients for whom an MBS Item was claimed than there are regular patients.

When considering the '12 months up to census date' we have 2 dates:

- LastClaimed - which is shared by all claims of the same type used for those items asking for the Days since last claim.
- DateLodged - which is specific to individual claims used to find out when the claim is done.

Both dates are used in the extract.

PI05 – HbA1c Recorded (Type 2 Diabetes Patients)

Description

Proportion of regular patients who are Indigenous, have type 2 diabetes and who have had an HbA1c (glycosylated haemoglobin) measurement result recorded at your primary health-care organisation within the previous 6 months **AND** the proportion of regular patients who are Indigenous, have type 2 diabetes and who have had an HbA1c measurement result recorded at your primary health-care organisation within the previous 12 months.

Business Rules:

- Patients with a history of Type 2 Diabetes are defined as having a condition listed under 'Past History' with one of the below DOCL codes:

diabm@niddm	Diabetes Mellitus - NIDDM
diabm@niddm	Diabetes Mellitus - Type II
diabm@niddm	NIDDM
diabm@niddm	NIDDM (Non-Insulin dependent diabetes mellitus)
diabm@niddm	Non-insulin dependent diabetes mellitus
diabm@niddm&rx%insu	Diabetes Type II requiring insulin
diabm@niddm&rx%insu	NIDDM - requiring insulin
- The latest HbA1c measurement must have been recorded in the previous 6 or 12 months. The minimum value for this calculation is 0 days. This is recorded through the 'Diabetes Assessment' screen or as a pathology result.
- HbA1c recorded is defined as patients whose records contain at least one of the following:
 - Measurement through Diabetes Assessment.
 - Pathology results containing one of the following LOINC codes: '17856-6','4548-4','59261-8','17855-8','4549-2','62388-4','53553-4' or the local codes (LOINC): 'C182','E482','21705226','30533253','31008465','31008445'
 - Pathology results containing the test names: '%GLYCATED HAEM%', '%GLYCOSYLATED HAEM%', '%HBA1C (GLYCATED HB%','Blood haemoglobin A 1 c', 'IFCCHbA1cPercent', 'IFCCHbA1c', 'IFCC HbA1c', 'B Glycosylated Haemoglobin (Hb A%', 'GLYCATED HAEM%', 'Glycated Haemoglob%', 'Glycated Hemoglobin', 'GLYCATEDHB', 'GLYCATEDHB%', 'GLYCOSYLATED HAEM%', 'GLYCOSYLATEDHB', 'Haemoglobin A1c%', 'HAEMOGLOBINA1C%', '+HbA1c%', '+Hb A1c%', 'HbA1c%', 'Hb A1c%', 'Hb A1C', 'HBA1C', 'HbA1c (IFCC)', 'HBA1C (SI)', 'HbA1c(IFCC)', 'HBA1C(SI)', 'HBA1CFRACTION', 'IFCC HbA1c', 'SI Hb A1C', 'HBA(HPLC)', 'Haemoglobin A1c%HBA1C(Glycated Haemoglob%', 'A1C', 'A1CFC ')

Calculation 1: HbA1c Measurement in the 6 Months Up to The Census Date

Numerator

Number of regular Indigenous patients who have had an HbA1c measurement result in the past 6 months.

Denominator

Number of regular Indigenous patients with type 2 diabetes as recorded in their file as per the below:

- Has Diabetes Type 2 as a 'reason for visit' OR
- Type Diabetes 2 Conditions are listed in the patient's history.

Calculation 2: HbA1c Measurement in the 12 Months Up to The Census Date

Numerator

Number of regular Indigenous patients who have had an HbA1c measurement result in the past 12 months.

Denominator

Number of regular Indigenous patients with type 2 diabetes as recorded in their file as per the below:

- Has Diabetes Type 2 as a 'reason for visit' OR
- Type Diabetes 2 Conditions are listed in the patient's history.

Reporting by Age Group

	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Records Excluded

Patients with type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose or impaired glucose tolerance.


Example - Add HbA1c Result from An Existing hl7 File

Please refer to this Online Help Guide: [The Holding File](#).

Example – How to Record HbA1c Value to Patient File

1. Open a patient's record.

2. Select **Clinical > Diabetes Record**.



3. The **Diabetes Follow Up** window appears.

Diabetes Follow Up

Assessments performed:

Assessment date

Last review by:

Ophthalmologist:

14/03/2019

Podiatrist:

14/03/2019

Dietitian:

14/03/2019

Endocrinologist:

14/03/2019

Diabetes educator:

14/03/2019

Last provided HypoKit:

14/03/2019

Major parameters:

Date	f Blood Gluc.	Height (cm)	Weight (kg)	HbA1c mmol/mol (%)	Creatinine	Pr

Assessment

Add Values

Graph

Close

- Click **Add Values**. The Diabetes Record window appears.

Diabetes record

Date:

Parameters				Urinalysis	
Weight (kg)		Height (cm)		Protein	
Systolic BP		Diastolic BP		Blood	
f Blood Gluc.		HbA1c (mmol/mol)		Glucose	
Total Chol.		HDL		Ketones	
Triglycerides		Creatinine		Leucocytes	
				Bilirubin	

Last review by:

Ophthalmologist	14/03/2019	Podiatrist	14/03/2019
Dietitian	14/03/2019	Endocrinologist	14/03/2019
Diabetes educator	14/03/2019		

Last Provided HypoKit:

- Input a value for HbA1c (either % or mmol/mol).
- Click **Save**.

Example - Record Type II Diabetes to Patient File in Patient History

- Select the **Past History** tab.
- Right click within the white space to add a new item.

Reactions: Smoking Hx: IHI No:


Warnings: MyHealthRecord:

Summary ☒ Current Rx ☐ Progress ☐ Past history ☐ Results ☐ Letters ☐ Documents ☐ Old script ☐

Year	Date	Condition	Side	Status	Summary
Edit item New item Delete item Print					

- The **New History Item** window appears.

4. Fill In the appropriate date and enter **Diabetes Mellitus - Type II** In the condition field.



5. Click **OK**.

Check

A regular patient who has had a test within the last 6 months should be counted in both the 6-month and 12-month calculations. For example, a regular patient who had a test 5 months ago would be counted in both.

For each age and sex at birth group:

- You cannot have more regular patients who have had HbA1c tests than the total number of regular patients with type 2 diabetes.
- You cannot have more regular patients who have had HbA1c tests in the previous 6 months than the number of regular patients who have had HbA1c tests in the previous 12 months.
- The total number of regular patients with type 2 diabetes should be the same for both the previous 6 months and the previous 12 months for Calculations 1 and 2.
- The total number of regular patients with type 2 diabetes should be the same as the total number of regular patients with type 2 diabetes in each relevant age and sex at birth group in these nKPIs: PI07, PI08, PI15, PI18 and PI23.
- The number of regular patients with type 2 diabetes should not exceed the number of regular patients in each group in these nKPIs: PI03 (adults), PI09 and PI16.

PI06 – Hba1c Results (Type 2 Diabetes Patients)

Description

Proportion of regular patients who are Indigenous, have type 2 diabetes and whose HbA1c measurement result was categorised as one of the following:

- Less than or equal to 7% (less than or equal to 53 mmol/mol).
- Greater than 7% but less than or equal to 8% (greater than 53 mmol/mol but less than or equal to 64 mmol/mol).
- Greater than 8% but less than 10% (greater than 64 mmol/mol but less than 86 mmol/mol).
- Greater than or equal to 10% (greater than or equal to 86 mmol/mol).

As recorded in the previous 6 months, **AND** as recorded in the previous 12 months.

Business Rules

- 'Type 2 Diabetes' as defined in PI05.
- The patient's HbA1c % is taken from the most recently recorded measurement or pathology result. The minimum value for days since HbA1c recorded is 0.
- HbA1c recorded as defined in PI05.

Calculation 1: Less Than or Equal To 7% In The 6 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement **less than or equal to 7%** (less than or equal to 53 mmol/mol) in the **6 months up to the census date**.

Recorded HbA1c matches the following boundary:

- HbA1c $\leq 7\%$ (≤ 53 mmol/mol).

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months** and fulfils the below criteria.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 6 months (Calculated in PI05).

Calculation 2: Less Than or Equal To 7% In The 12 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement **less than or equal to 7%** (less than or equal to 53 mmol/mol) in the **12 months up to the census date**.

Recorded HbA1c% matches the following boundary:

- HbA1c $\leq 7\%$ (≤ 53 mmol/mol).

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 12 months (Calculated in PI05).

Calculation 3: More Than 7% But Less Than or Equal To 8% In The 6 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement **more than 7% but less than or equal to 8%** (greater than 53 mmol/mol but less than or equal to 64 mmol/mol) in the **6 months up to the census date**.

Recorded HbA1c% matches the following boundary:

- HbA1c >7% and <=8% (>53 but <= 64 mmol/mol)

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 6 months Calculated in PI05).

Calculation 4: More Than 7% But Less Than or Equal To 8% In The 6 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement **more than 7% but less than or equal to 8%** (greater than 53 mmol/mol but less than or equal to 64 mmol/mol) in the **12 months up to the census date**.

Recorded HbA1c% matches the following boundary:

- HbA1c >7% and <=8% (>53 but <= 64 mmol/mol).

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 6 months (Calculated in PI05).

Calculation 5: More Than 8% But Less Than 10% In The 6 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement **more than 8% but less than 10%** (greater than 64 mmol/mol but less than 86 mmol/mol) in the **6 months up to the census date**.

Recorded hba1cPercent matches the following boundary:

- HbA1c >8% and <10% (>64 but <= 86 mmol/mol)

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 6 months (Calculated in PI05).

Calculation 6: More Than 8% But Less Than 10% In The 12 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement **more than 8% but less than 10%** (greater than 64 mmol/mol but less than 86 mmol/mol) in the **12 months up to the census date**.

Recorded Hba1c% matches the following boundary:

- HbA1c >8% and <10% (>64 but <86 mmol/mol).

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 12 months (Calculated in PI05).

Calculation 7: More Than 10% In The 6 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement of **10% or more** (greater than or equal to 86 mmol/mol) in the **6 months up to the census date**.

Recorded HbA1c% matches the following boundary:

- HbA1c >=10% (>=86 mmol/mol).

Denominator

The number of regular Indigenous patients have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 6 months (Calculated in PI05).

Calculation 8: More Than 10% In The 12 Months Up to The Census Date

Numerator

The number of regular Indigenous patients that have had an HbA1c measurement of **10% or more** (greater than or equal to 86 mmol/mol) in the **12 months up to the census date**.

Recorded HbA1c% matches the following boundary:

- HbA1c $\geq 10\%$ (≥ 86 mmol/mol).

Denominator

The number of regular Indigenous patients that have had an HbA1c measurement result recorded in the **same 6 months**.

- Has Diabetes (See PI05) AND
- Has a record of HbA1c within 12 months (Calculated in PI05).

Reported by Age Group

	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Records Excluded

- Patients with type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose or impaired glucose tolerance were excluded.

Notes

- Only the most recently recorded result from an HbA1c test is used. This means that if a patient has had several tests, we only include the result from the most recent test.
- Results from all relevant pathology tests. If your organisation doesn't have a good system for adding pathology results to patient records, you will need to make sure they have been included in the correct field for all patients in this nKPI.

Example – Recording HbA1c Value to a Patient's File

HbA1c value may be recorded following the method outlined in [PI05](#).

Example - Record Type 2 Diabetes to a Patient's File

History of diabetes may be recorded following the method outlined in [PI05](#).

Check

A regular patient who has had a test within the last 6 months should be counted in both the 6-month and 12-month calculations. For example, a regular patient who had a test 5 months ago would be counted in both.

For each age and sex at birth group:

- You cannot have more people with HbA1c test results in each group than the total number of regular patients who've had HbA1c tests recorded.

PI07 – Chronic Disease Management Plan prepared

Description

Proportion of Indigenous regular clients who have a chronic disease (Type 2 diabetes) and for whom a Chronic Disease Management Plan was prepared within the previous 24 months as indicated by:

An MBS-rebate Chronic Disease Management Plan:

- In-person MBS-rebate items:
 - Historical: 229, 721
 - New (as of 1 July 2025): 65, 392, 967, 3939
- Telehealth MBS-rebate items:
 - Historical: 92024, 92055
 - New (as of 1 July 2025): 92029, 92030, 92060, 92061

Note: This indicator will report on both historical and new item codes during the transitional period from 1 July 2025 to 30 June 2027.

At this stage, Type 2 Diabetes is the only chronic disease included for this indicator.

Business Rules

- 'Type 2 Diabetes' as defined in PI05.
- 'MBS Item' claimed as defined in PI03.

Calculation 1: Indigenous regular clients with a chronic disease (Type 2 diabetes) for whom a GP Management Plan (MBS items 721, 229, 965, 392, 967, 393) was claimed within the previous 24 months

Numerator

Number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]** who have Type 2 diabetes and for whom an in-person MBS-rebate Chronic Disease Management Plan (MBS items: 721, 229, 965, 392, 967, 393) was prepared within the previous 24 months. (MBS Claimed explained in PI03).

Denominator

Total number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]** who have Type 2 diabetes.

Calculation 2: Indigenous regular clients with a chronic disease (Type 2 diabetes) for whom a GP Management Plan (MBS items 92024, 92055, 92060, 92030, 92061) was claimed within the previous 24 months

Numerator

Number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]** who have Type 2 diabetes and for whom an in-person MBS-rebate Chronic Disease Management Plan (MBS items: 92024, 92055, 92060, 92030, 92061) was prepared within the previous 24 months. (MBS Claimed explained in PI03).

Denominator

Total number of **[Disaggregation: Sex at Birth]** Indigenous regular clients aged **[Disaggregation: Age]** who have Type 2 diabetes.

Reporting by Age Group

	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Records Excluded

- Patients with type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose or impaired glucose tolerance.
- Do not include patients who have only had a GP Management Plan review (Medicare Item 732) within the recording period.

Notes

Include in the Numerator only those patients whose GP Management Plan was claimed by your organisation under MBS Item 721.

- Services taking part in the Health Care Homes Trial: For the duration of the HealthCare Homes trial (currently 1 October 2017 to 30 November 2019), patients who are part of the trial will be deemed to have had an MBS Item 721 claimed if there is evidence of a GP Management Plan recorded.
- While Type 2 Diabetes is the only chronic disease needed for this nKPI, MedicalDirector Insights also reports on: Cardiovascular disease, Chronic Obstructive Pulmonary Disease and Chronic Kidney Disease.

Example – Creating and Processing a 721 Claim

The method for creating and processing a 721 claim can be seen in PI03.

Example – Recording History of Type 2 Diabetes

A history of Type 2 Diabetes may be added to the patient file by following the method outlined in PI05.

Check

For each age and sex at birth group:

- You cannot have more patients for whom an MBS Item (for managing type 2 diabetes) was claimed than there are patients with type 2 diabetes.
- The total number of regular patients with type 2 diabetes should be the same as the total number of regular patients with type 2 diabetes in each relevant age and sex at birth group in these nKPIs: PI05, PI08, PI15, PI18 and PI23.
- The total number of regular patients with type 2 diabetes should not exceed the corresponding number of total regular patients for corresponding groups in these nKPIs: PI03 (adults), PI09 and PI16.

PI09 – Smoking Status Recorded

Description

Proportion of regular patients who are Indigenous, aged 11 and over **and** whose smoking status has been recorded at your primary health-care organisation within the previous 24 months.

Business Rules

Patient's smoking status is recorded in the smoking tab in the patient's details menu.

Numerator

The number of regular Indigenous patients aged over 11 that have had their smoking status recorded in the **24 months up to the census date**.

Denominator

The number of your regular Indigenous patients in each age and sex at birth group.

Reporting by Age Group

	Age (years)						
	11 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓

Notes

- Where an Indigenous regular patient's tobacco smoking status does not have an assessment date assigned within the Patient Information Record System (PIRS), smoking status as recorded in the PIRS should be treated as current (that is, as having been updated within the previous 24 months).
- Results arising from measurements conducted outside of the organisation that are known by the organisation should be included.


Check

For each age and sex at birth group:


- You cannot have more people with their smoking status recorded than you have patients of the same age and sex at birth.
- The number of regular patients with smoking status recorded should be the same as the total number of regular patients of the same age and sex at birth with their smoking status recorded in this nKPI: PI10.
- The total number of regular patients should be the same as the total number of patients of the same age and sex at birth in these nKPIs: PI03 (adults) and PI16.

Example – Record Smoking Status

1. Open a patient's record.
2. Select **Patient > Details**.



3. Select the **Smoking** tab.



The screenshot shows the 'Patient Details' dialog box with the 'Smoking' tab selected. The 'Smoker' dropdown is set to 'Never smoked'. Other fields include 'Date of assessment' (14/03/2019), 'Frequency', 'Number of cigarettes' (5), 'Year commenced' (1980), 'Duration' (39yrs), 'Stage of change assessment', 'Last quit attempt' (14/03/2019), and 'Duration of longest period of abstinence'. There are also checkboxes for 'Smoking cessation intervention discussed with patient', 'Update address for all family members', and 'Auto-capitalise names'. Buttons for 'View Patient Education Leaflet', 'Reference', 'Save', and 'Cancel' are visible.

4. Set smoking status via the **Smoker** dropdown.
5. Click **Save**.

Check

For each age and sex at birth group:

- You cannot have more people with their smoking status recorded than you have clients of the same age and sex at birth.
- The number of regular clients with smoking status recorded should be the same as the total number of regular clients of the same age and sex at birth with their smoking status recorded in this nKPI: PI10.
- The total number of regular clients should be the same as the total number of clients of the same age and sex at birth in these nKPIs: PI03 (adults) and PI16.

PI10 – Smoking Status Result

Description

Proportion of regular patients who are Indigenous, aged 11 and over **and** whose smoking status has been recorded within the previous 24 months as one of the following:

- Current smoker.
- Ex-smoker.
- Never smoked.

There's no agreement on how long a person needs to have quit smoking to be considered an ex-smoker rather than a smoker—what is put on the record is a clinical judgement. To be counted as having ever smoked, the person must have smoked more than **100 cigarettes in total (or equivalent)**.

Business Rules

'Smoking status' as defined in PI09.

Calculation 1: Current Smokers

Numerator

The number of regular Indigenous patients that are entered as **current smokers** in the **24 months up to the census date**.

- All patients who are 'daily smoker', 'weekly smoker' and 'irregular smoker'—add them together and count them as 'current smoker'.

Denominator

The number of regular Indigenous patients that have their smoking status recorded in the **same 24 months**.

Calculation 2: Ex-smokers

Numerator

The number of regular Indigenous patients that are entered as **ex-smokers** in the **24 months up to the census date**.

Denominator

The number of regular Indigenous patients that have their smoking status recorded in the **same 24 months**.

Calculation 3: Non-smokers

Numerator

The number of regular Indigenous patients that are entered as **never smoked** in the **24 months up to the census date**.

Denominator

The number of regular Indigenous patients that have their smoking status recorded in the **same 24 months**.

Reporting by Age Group

	Age (years)						
	11 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓

Notes

Where an Indigenous regular patient's tobacco smoking status does not have an assessment date assigned within the Patient Information Record System (PIRS), smoking status as recorded in the PIRS should be treated as current (that is, as having been updated within the previous 24 months).

Example - Record Smoking Status

See example listed in PI09

Check

For each age and sex at birth group:

- You cannot have more people with a smoking status (sum of current smokers, ex-smokers and those who never smoked) than you have patients of the same age and sex at birth with any smoking status in their record.
- The total number of regular patients with their smoking status recorded should be the same as the number of regular patients of the same age and sex at birth with their smoking status recorded in PI09.

PI11 – Smoking During Pregnancy

Description

Proportion of female regular clients who are Indigenous, who gave birth within the previous 12 months and whose smoking status was recorded during pregnancy as one of the following:

- Current smoker.
- Ex-smoker.
- Never smoked.

There's no agreement on how long a person needs to have quit smoking to be considered an ex-smoker rather than a smoker—what is put on the record is a clinical judgement. To be counted as having ever smoked, the person must have smoked more than **100 cigarettes in total (or equivalent)**.

Business Rules

- Counts clients with a recorded smoking status, who recorded 3 or more visits within the previous 2 years.
- Includes live births and still births if the birthweight is at least 400 grams OR the gestational age was 20 weeks or more.
- When the birthweight is not recorded, then the birthweight is considered as < 400 grams.
- Include only the most recent smoking status recorded prior to the completion of the latest pregnancy. Where a smoking status does not have an assessment date assigned within the CIS, smoking status should not be counted.
- Include results arising from measurements conducted outside of the organisation that are known by the organisation.

Calculation 1: Current Smoker

Numerator

How many regular Indigenous female patients aged **[Disaggregation: Age]** who gave birth within the previous 12 months whose most recent smoking status has been recorded as **current smoker**

- All patients who are 'daily smoker', 'weekly smoker' and 'irregular smoker'—add them together and count them as 'current smoker'.
- Note: If smoking status has been recorded, we do not have a date limit on when this data point was recorded.

Denominator

The number of regular Indigenous female patients aged **[Disaggregation: Age]** gave birth within the previous 12 months who had their smoking status recorded in the **same 12 months**.

Calculation 2: Ex-smoker

Numerator

How many regular Indigenous female patients aged **[Disaggregation: Age]** who gave birth within the previous 12 months whose most recent smoking status has been recorded as **ex-smoker**.

Note: Provided smoking status has been recorded we do not have a date limit on when this data point was recorded.

Denominator

The number of regular Indigenous female patients aged **[Disaggregation: Age]** gave birth within the previous 12 months who had their smoking status recorded in the **same 12 months**.

Calculation 3: Non-smoker

Numerator

How many regular Indigenous female patients aged **[Disaggregation: Age]** who gave birth within the previous 12 months whose most recent smoking status has been recorded as **never smoked**.

Note: Provided smoking status has been recorded we do not have a date limit on when this data point was recorded.

Denominator

The number of regular Indigenous female patients aged **[Disaggregation: Age]** gave birth within the previous 12 months who had their smoking status recorded in the **same 12 months**.

Reporting by Age Group

	Age (years)		
	<20 years	20 to <35	35+
Female	✓	✓	✓


Notes

- Live births and stillbirths if the birthweight was at least 400 grams or the gestational age was 20 weeks or more.
- Where an Indigenous regular patient's tobacco smoking status does not have an assessment date assigned within the Patient Information Record System (PIRS), smoking status as recorded in the PIRS should be treated as current (that is, as having been updated within the previous 12 months).

Example - Recording Pregnancy


1. Open a female patient's record.
2. Select the **Obstetric** tab.

- Right-click in the white space to add a **New Item**.




The screenshot shows the MedicalDirector software interface. The patient record for Mrs. Jennifer ANDREWS (48yrs 10mths) is displayed. The 'Obstetric' tab is selected, and a right-click context menu is open over it, showing options: 'Edit item', 'New item' (highlighted), 'Delete item', and 'Print'.

- Input the value for **Date of LMP** or gestational age by scan and click **Save**.



The screenshot shows the 'Gestational Age Calculator' dialog box. The 'Date of LMP' is set to 14/03/2019. The 'By LMP' radio button is selected. The 'Calculate' button is highlighted.




The screenshot shows the 'Gestational Age Calculator' dialog box with the results of the calculation. The 'Date of Scan' is 14/03/2019. The 'By Scan' radio button is selected. The 'Calculate' button is highlighted. The results show: E.D.C. is 25/11/2019, Current gestational age is 3 weeks, and 3 days.

Week	Start date
1	25/02/2019
2	04/03/2019
3	11/03/2019
4	18/03/2019


Example - Recording Pregnancy Result

1. Click the **End Pregnancy** button.




The screenshot shows the MedicalDirector interface with a toolbar at the top containing icons for Summary, Rx, Current Rx, Progress, Past history, Results, Letters, Documents, Old scripts, Imm., and Cervix. Below the toolbar is a navigation bar with buttons: G1 P0 C, Bl.Gp. B+, New pregnancy, Record visit, End pregnancy (highlighted with a yellow box), Past pregnancy, Path. results, and Parity. Below the navigation bar is a table with columns: Pregnancy, LNMP, EDC, EDC By Scan, Outcome, Date, Gest., B.Weight, and Name. The table contains one row with the following data: 1, 18/02/2019, 25/11/2019, 0.

2. When the **Antenatal Record** dialogue box appears, click **Yes**.



The screenshot shows the 'Antenatal record' dialogue box. It contains a red 'X' icon and the text: 'Ending this pregnancy will prevent any further antenatal visits being recorded! Are you sure that you wish to proceed?'. At the bottom, there are two buttons: 'Yes' (highlighted with a blue box) and 'No'.

3. When the **Past Obstetric History** dialogue box appears:
 - Record a **Finish Date**.
 - Select an **Outcome** from dropdown list (e.g., Normal vaginal delivery).
 - Enter Baby Name, Baby Weight, Baby Sex at Birth.




The screenshot shows the 'Past Obstetric History' dialogue box. It contains the following fields and options:

- Pregnancy No.:** 5
- Finish date:** 14/11/2016 (selected from a dropdown menu)
- Gestation (wks):** (empty field)
- Outcome:** (dropdown menu with 'Para' selected, and radio buttons for 'Termination' and 'Miscarriage')
- Notes:** (text area)
- Labour:**
 - Stage 1:** (empty field)
 - Stage 2:** (empty field)
 - Stage 3:** (empty field)
 - Episiotomy/tear:** (empty field)
- Baby:**
 - Name:** (empty field)
 - Birth weight (gms):** (empty field)
 - Gender:** (dropdown menu)
 - Feeding:** (radio buttons for 'Breast' and 'Bottle')
 - Apgar 1:** (empty field)
 - Apgar 5:** (empty field)

At the bottom, there are two buttons: 'Save' and 'Cancel'.

4. Click **Save**.
5. When the **End Pregnancy** dialog box appears, select **No** and do not create a patient file.



Example - Record Smoking Status

Smoking status should be recorded as outlined in PI09

Check

For each age group (<20, 20 to <35 and 35 and over):

- You cannot have more people with a smoking status (sum of current smokers, ex-smokers and who never smoked) than you have patients of the same age with any smoking status in their record.
- The number of regular patients who gave birth within the previous 12 months with their smoking status recorded should not be more than the total number of regular patients of the same age and sex at birth with their smoking status recorded in PI09.
- The total number of regular patients who gave birth within the previous 12 months with their smoking status recorded should be equal to or less than the total number of regular patients who gave birth in the previous 12 months in the relevant age groups in PI13 as per the table below:

PI11	PI13
<20	<20
20 to <35	20 to <35
35+	35+

PI12 – Body Mass Index

Description

Proportion of regular clients who are Indigenous, aged 18 and over who have had their body mass index (BMI) classified as:

- underweight (<18.50).
- normal weight (≥ 18.50 but <25).
- overweight (≥ 25 but <30).
- obese (≥ 30).
- not calculated.

Within the previous 24 months.

Business Rules

- Only the **most recent** BMI is considered.
- Only weight measurements recorded in the last 24 months are considered.
- Only height measurements recorded since the patient turned 18 are considered.
- If any of the above conditions are not met, then the patient is excluded from the calculation.

Calculation 1: Underweight

Numerator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age] who have had their BMI classified as Underweight (BMI <18.50) within the previous 24 months.

Denominator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age].

Calculation 2: Normal weight

Numerator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age] who have had their BMI classified as Normal weight (BMI ≥ 18.50 AND <25) within the previous 24 months

Denominator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age].

Calculation 3: Overweight

Numerator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age] who have had their BMI classified as Overweight (BMI ≥ 25 and <30) within the previous 24 months.

Denominator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age].

Calculation 4: Obese

Numerator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age] who have had their BMI classified as Obese (BMI ≥ 30) within the previous 24 months.

Denominator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age].

Calculation 5: Not calculated

Numerator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age] who have not had their BMI classified within the previous 24 months.

Denominator

The number of indigenous regular [Disaggregation: Sex at Birth] clients aged [Disaggregation: Age].

Disaggregation

A visual representation of the disaggregation per calculation (overweight, obese, normal weight, underweight and not calculated) can be seen below:

	Age (years)					
	18 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓
Male	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓
Male	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓

Notes

- 'Body Mass Index (BMI)' – is a measure of an adult's weight (body mass) relative to height used to assess the extent of weight deficit or excess where height and weight have been measured. BMI is the weight in kilograms divided by height in metres squared (WHO 2000).
- 'Constituent Elements of a BMI' – weight in kilograms, recorded within the previous 24 months; height in meters, recorded since the client turned 18 years of age.
- BMI is more likely to be recorded for certain client groups, such as those with diabetes.
- BMI is more likely to be recorded if a client appears underweight, overweight, or obese. This could result in a higher apparent proportion of underweight, overweight, or obese patients.
- If height and weight are recorded for a client, their BMI is also considered as recorded.

Example – How to Enter BMI


Use height and weight measured based on the following rules:

- A height taken since the client turned 18.
- A weight taken within the previous 24 months.


The BMI needs to be recorded in the Weight Calculator per the below steps:

1. Open a patient's record.

2. Select **Tools > Tool Box > Weight**.



3. Input height and weight values.



Weight Calculator

Date: 7/08/2017 Time: 1:49:18 PM Gender: Female Age: 47 Height: 175 Patient ID: 31

Blood Glucose Blood Pressure CV Risk ECG INR Renal Function Respiratory **Weight**

Current Measurements View: All

Height: 175 (cm)
 Weight: 64 (kg)
 BMI: 20.9
 Waist: (cm)
 Hip: (cm)
 Waist/Hip Ratio:
 Healthy weight range = 56.7 - 76.6 Kg
 Median range value = 61.3 Kg

Date	Time	Type	Value
04/11/2003	13:50:08	Weight	65
04/11/2003	13:50:08	BMI	21.2
10/10/2006	09:55:35	Height	175
10/10/2006	09:55:35	Weight	65
10/10/2006	09:55:35	BMI	21.2
05/12/2006	10:28:49	Weight	64
03/01/2007	10:35:50	Height	175
03/01/2007	10:35:50	Weight	68
03/01/2007	10:35:50	BMI	22.2
22/03/2007	10:44:12	Weight	69
10/05/2007	10:47:51	Weight	72.0
30/08/2007	10:52:44	Height	175
30/08/2007	10:52:44	Weight	65

Clear Print Reference Edit View Save Close

4. Click **Save**.

Check:

For each age and sex at birth group:

- You cannot have more overweight or obese or normal weight or underweight or not calculated regular clients than you have regular clients whose BMI has been recorded.
- The number of regular clients who have had their BMI recorded must be less than in the total number of regular clients in the relevant age and sex at birth groups in these nKPIs: PI03 (adults), PI09 and PI16. The only exception is if you have very few clients in the age or sex at birth group.

PI13 – First Antenatal Care Visit

Description

Proportion of female Indigenous regular patients, who gave birth within the previous 12 months and who had gestational age recorded at their first antenatal care visit as one of the following:

- Less than 11/40 weeks.
- At or after 11/40 and less than 14/40.
- At or after 14/40 weeks and less than 20/40 weeks.
- At or after 20/40 weeks.
- No result.
- Did not attend an antenatal care visit.

Business Rules

'Parity Birth' and 'Gestational Age' is defined as per [PI11](#).

- When the birthweight is not recorded, then the birthweight is considered as < 400 grams.

Calculation 1: Less Than 11/40 Weeks

Numerator: The number of female Indigenous regular patients aged *[Disaggregation: Age]* who gave birth within the previous 12 months, who had gestational age recorded at their first antenatal care visit with results less than 11/40 weeks of gestation.

Denominator: The number of female Indigenous regular patients aged *[Disaggregation: Age]* who gave birth within the previous 12 months.

Calculation 2: At or after 11/40 and less than 14/40 Weeks

Numerator: The number of female Indigenous regular patients aged *[Disaggregation: Age]* who gave birth within the previous 12 months, who had gestational age recorded at their first antenatal care visit with results at or after 11/40 weeks and less than 14/40 weeks of gestation.

Denominator: The number of female Indigenous regular patients aged *[Disaggregation: Age]* who gave birth within the previous 12 months.

Calculation 3: At or after 14/40 weeks and less than 20/40 Weeks

Numerator: The number of female Indigenous regular patients aged *[Disaggregation: Age]* who gave birth within the previous 12 months, who had gestational age recorded at their first antenatal care visit with results at or after 14/40 weeks to less than 20/40 weeks of gestation.

Denominator: The number of female Indigenous regular patients aged *[Disaggregation: Age]* who gave birth within the previous 12 months.

Calculation 4: At or after 20/40 Weeks

Numerator

The number of female Indigenous regular patients aged [Disaggregation: Age] who gave birth within the previous 12 months, who had gestational age recorded at their first antenatal care visit with results at or after 20/40 weeks of gestation.

Denominator

The number of female Indigenous regular patients aged **[Disaggregation: Age]** who gave birth within the previous 12 months.

Calculation 5: No Gestational Age Recorded

Numerator

The number of female Indigenous regular patients aged [Disaggregation: Age] who gave birth within the previous 12 months, who had no gestational age recorded at their first antenatal care visit.

Denominator

The number of female Indigenous regular patients aged **[Disaggregation: Age]** who gave birth within the previous 12 months.

Calculation 6: No Antenatal Visit Recorded

Numerator

The number of female Indigenous regular patients aged [Disaggregation: Age] who gave birth within the previous 12 months, who had no antenatal care visit recorded.

Denominator

The number of female Indigenous regular patients aged **[Disaggregation: Age]** who gave birth within the previous 12 months.

Reporting by Age Group

Mother's Age (years)		
<20	20 to <35	35 and over
✓	✓	✓
✓	✓	✓

Notes


- Include live births and still births if the birthweight was at least 400 grams or the gestational age was 20 weeks or more.
- When the birthweight is not recorded, then the birthweight is considered as < 400 grams

Example - Add New Pregnancy Record

A new pregnancy can be recorded following the method outlined in PI11


Example - Record First Antenatal Visit

1. Select a pregnancy record from the list.
2. Click **Record Visit**.



Pregnancy	LNMP	EDC	EDC By Scan	Outcome	Date	Gest.	B.Weight	Name
1	02/01/2019	09/10/2019				0		

3. Input the date as the visit date.



Antenatal record

Date: 14/03/2019 Weight:

B.P. Systolic: Diastolic:

Oedema: Urine:

Size (weeks) Calc: 10 Clin:

Fundal height (cm): Foetal heart:

Presentation:

Comments:

Save Cancel

4. Click **Save**.

Check

For each age group:

- You cannot have more mothers with gestational age recorded at their first antenatal visit than number of mothers who have given birth.
- You cannot have more mothers with no gestational age recorded at their first antenatal visit than number of mothers who have given birth.
- The number of mothers with any gestational age recorded, the number of mothers without gestational age recorded, and the number of mothers who did not have an antenatal visit should add up to the total number of mothers who've given birth. (Make a note in the submission comments if this is not the case and there is a reason for the difference).

PI14 – Influenza Immunisation

Description

Proportion of Indigenous regular clients aged 6 months and over who are immunised against influenza.

Business Rules

Patient's Influenza Immunisation status is defined in the 'Immunisation record' under the code 'flu' or description 'Influenzae'. If this is present in the patient's file, the date between the immunisation date and date of reporting is calculated. If this date difference is ≥ 12 months, the record is included.

Flu vaccines included are listed below:

Vaccine	Vaccine code	Vaccine	Vaccine code
Afluria Quad	AFLR, AFLQUA, QUADAF	Fluzone High-Dose Quadrivalent	FLHDQD
Agrippal	AGRPAL	Generic Influenza	GNFLU
Fluad	FLUAD	Influvac	INFLUV
Fluad Quad	FLUQAD	Influvac Tetra	INFLTA
Fluarix Tetra	FLUTET	Panvax H1N1	PANVAX
Flucelvax Quad	FCELQD	Vaxigrip	VAXGRP
FluQuadri	FQUAD	Vaxigrip Junior	VGRJNR
FluQuadri Junior	FQDJN	Vaxigrip Tetra	VAXTET
Fluvax	FLUVAX	Influvac Junior	
Fluvax Junior	FVXJNR	Influenza Vaccine, Tetravalent	
Fluvirin	FLVRN	Intanza 9MCG	
Fluzone	FLUZNE	FluMist	FLUMIS
Fluzone High-Dose	FLUHID	Pandemic (H1N1) 2009 Influenza	

Numerator

The number of [Disaggregation: Sex at Birth] Indigenous regular clients aged [Disaggregation: Age] who are immunised against influenza in the 12 months up to the census date.

Note: These need to be recorded in the immunisations tab like PI04.

Denominator

The number of [Disaggregation: Sex at Birth] Indigenous regular clients aged [Disaggregation: Age].


Records Excluded

Patients in the Numerator if they have not been vaccinated, regardless of the reason (you will still include them in the Denominator). People who have not been vaccinated are at a higher risk of catching influenza.

Example - Record Influenza Immunisation


1. Open a patient's record.
2. Select the **Immunisation** tab.

- Right-click in the white space to add a **New Item**.



Reactions: [dropdown] Smoking Hx: 7 Smoker IHI No: [dropdown]
 Warnings: Pregnant: 10 weeks, 1 days, EDC = 09/10/2019 MyHealthRecord: [dropdown]
 Summary Rx Current Rx Progress Past history Results Letters Documents Old scripts **Imm.** Cervical Sc
 Date Immunisation Sequence Batch Consent Provided by Vaccinator Site
 Edit item
 New item
 Delete item
 Print
 Refresh

- Enter date, type (Influenza) site and sequence.



Vaccination Window
☐ Vaccine given elsewhere
 Vaccinator: Dr A Practitioner
 Consent provided by: Patient ☐ Vaccination declined
 Date: 14/03/2019
 Type: INFLUENZA
 Site: Deltoid SC, Left side
 Sequence: 2
 Batch No: 1 ☐ Store batch No
 Comment:
 [Text Area]
 Mark for recall Save Cancel

- Click **Save**.

Check

For your regular patients in the age groups:

- You cannot have more patients being immunised than the total number of patients who fall under that age category.
- You cannot have more men being immunised than the total number of men who fall under that age category.
- You cannot have more women being immunised than the total number of women who fall under that age category.

	Age							
	6 months to <5 years	5 to <15 years	15 to <25 years	25 to <35 years	35 to <45 years	45 to <55 years	55 to <65 years	65 years+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

PI16 – Alcohol Consumption Recorded

Description

Proportion of regular patients who are Indigenous, aged 15 and over and who have had their alcohol consumption status recorded at your primary health-care organisation within the previous 24 months.

Business Rules

Alcohol consumption status recorded is defined as patients for whom an Audit-C measurement exists.

Numerator

The number of regular Indigenous patients that have had their alcohol consumption status recorded in the 24 months up to the census date.

- Any record of alcohol consumption. This could include a record of:
 - Whether the regular patient consumes alcohol.
 - The amount and frequency of the regular patient's alcohol consumption.
 - The results of tests such as the AUDIT or AUDIT-C.
- Where an Indigenous regular patient's alcohol consumption status does not have an assessment date assigned in the Patient Information Record System (PIRS), alcohol consumption status as recorded in the PIRS should be treated as current (that is, as having been updated within the previous 24 months).

Denominator

The number of your regular Indigenous patients that were there in each age and sex at birth group.


Reporting by Age Groups

	Age (years)					
	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓


Example - Recording Alcohol Consumption

1. Open a patient's record.

2. Select **Assessment > Audit-C**.



3. The **Audit-C Assessment** is presented.



The screenshot shows the 'Patient Details' window with the 'Audit-C Assessment' tab selected. The form includes the following sections:

- Date of assessment:** 10/11/2016
- Audit-C Assessment:**
 - 1. How often do you have a drink containing alcohol?
 - ☐ Never
 - ☒ Monthly or less
 - ☐ 2-4 times a month
 - ☐ 2-3 times a week
 - ☐ 4 or more times a week
 - 2. How many standard drinks containing alcohol do you have on a typical day?
 - ☐ 1 or 2
 - ☐ 3 or 4
 - ☐ 5 or 6
 - ☐ 7 to 9
 - ☒ 10 or more
 - 3. How often do you have six or more drinks on one occasion?
 - ☐ Never
 - ☐ Less than monthly
 - ☐ Monthly
 - ☐ Weekly
 - ☒ Daily or almost daily
- Audit-C Total Score: 9**

In men a score of 4 or more and in women a score of 3 or more is considered positive, optimal for identifying hazardous drinking or active alcohol use disorders. The guidelines to reduce health risks from drinking alcohol provide further assessment and treatment options.

Patient concerned about drinking?

 - ☐ Yes
 - ☐ No
 - ☒ Don't know

Buttons: View Alcohol Guidelines, Reference, New Assessment

Currently displaying data from assessment performed on 10/11/2016. Click 'New Assessment' to conduct a new assessment.
- Table:**

Date	Time	Score	Concerns	Comments
10/11/2016	10:55:22	9	Don't know	No
10/11/2016	09:53:28	2	Don't know	No
- Comments:** (Text area)
- Buttons:** Delete, Save, Cancel
- Footer:**
 - ☐ Update address for all family members
 - ☐ Auto-capitalise names

4. Input data and click Save.

Check

For each age and sex at birth group:

- You cannot have more regular patients with a record of their alcohol consumption status than the total number of regular patients.
- The number of regular patients who had their alcohol consumption status recorded should be greater than or equal to the total number of regular patients who had an AUDIT-C recorded in this nKPI: PI17.
- The total number of regular patients should be the same as the total number of regular patients in these nKPIs: PI03 (adults) and PI09.

PI18 – Kidney Function Test Recorded (Type 2 Diabetes or CVD Patients)

Description

Proportion of regular clients who are Indigenous, who are recorded as having:

- Type 2 diabetes OR
- Cardiovascular disease (CVD) OR
- Type 2 diabetes and/or CVD

AND who have had either:

- An estimated glomerular filtration rate (eGFR) AND an albumin/creatinine ratio (ACR) test result recorded.
- Only an eGFR test result recorded.
- Only an ACR test result recorded.
- Neither an eGFR NOR an ACR test result recorded within the previous 12 months.

Business Rules

- Type 2 Diabetes recorded as outlined in PI05.
- History of CVD recorded is defined as patients with a condition listed as at least one of the docles mentioned in the DOCLE list (Appendix A):
- ACR or other micro albumin test result recorded is defined as patients whose records contain at least one of the following:
 - Pathology results containing one of the following LOINC codes: '14959-1', '32294-1', '30001-2', '30000-4', '9318-7'.
 - Pathology results containing the test names: 'ACR', 'Albumin/creatinine ratio urine', 'Alb/Creat', 'Albumin/Creat', 'Alb Creat Ratio', 'Alb/Cre', 'Albumin/Creat', 'Albumin/Creatinine', 'Albumin/Creatinine Ratio', 'Creatinine (Ref Lab)', 'MA Creat Ratio', 'Malb/Crea', 'Microalbumin Ratio', 'S Creatinine', 'Serum Creatinine', 'U Ab Creat Ratio', 'Urinary Albumin/Creat Ratio', 'Urinary Albumin/Creatinine Ratio', 'UrineAlbumin/CreatinineRatio', 'ALBCR', 'LBCR'
 - Electronic results and manually recorded ACR measurements of type: 'MALB', 'MALBUN' or 'MALDATE'.
- eGFR recorded is defined as patients whose records contain at least one of the following:
 - Pathology results containing one of the following LOINC codes: '33914-3', '62238-1', '50210-4', '98979-8', '98980-6'.
 - Pathology results containing the test names: 'eGFR', 'GFR estimated', 'Glomerular filtration rate estimated'
 - Pathology results containing one of the following UNITS: 'ml/min/1.73m2', 'ml/min/1.73 m2', 'ml/min/1.73sqm'
 - Electronic results and manually recorded eGFR measurement of type: 'EGFR'.

- HBA1C recorded is defined as patients whose records contain at least one of the following
 - Pathology results containing one of the following LOINC codes: '17856-6', '4548-4', '59261-8'.
 - Pathology results containing the test names: 'GLYCATED HAEM', 'GLYCOSYLATED HAEM', 'HBA1C (GLYCATED HB', 'Blood haemoglobin A 1 c', 'IFCCHbA1cPercent', 'IFCCHbA1c', 'Glycated Haemoglob', 'Glycated Hemoglobin', 'GLYCATEDHB', 'GLYCATEDHB', 'GLYCOSYLATEDHB', 'Haemoglobin A1c%HBA1C(Glycated Haemoglob%', 'HAEMOGLOBINA1C', 'Hb A1C', 'HBA(HPLC)', 'HBA1C', 'HbA1c (IFCC)', 'HBA1C (SI)', 'HbA1c(IFCC)', 'HBA1C(SI)', 'HBA1CFRACTION', 'IFCC HbA1c', 'SI Hb A1C', 'A1CIFC', 'A1C'

Calculation 1: Type 2 Diabetes

Numerator

The number of Indigenous regular [Disaggregation: Sex at Birth] clients with Type 2 Diabetes aged [Disaggregation: Age] who have had [Disaggregation: Tests] recorded within the previous 12 months.

Denominator

The number of Indigenous regular [Disaggregation: Sex at Birth] clients with Type 2 Diabetes aged [Disaggregation: Age] (see [PI05](#)).

Calculation 2: CVD

Numerator

The number of Indigenous regular [Disaggregation: Sex at Birth] clients with CVD aged [Disaggregation: Age] who have had [Disaggregation: Tests] recorded within the previous 12 months.

Denominator

The number of Indigenous regular [Disaggregation: Sex at Birth] clients with CVD aged [Disaggregation: Age] (see [PI05](#)).

Calculation 3: Type 2 Diabetes OR CVD

Numerator

The number of Indigenous regular [Disaggregation: Sex at Birth] clients with Type 2 Diabetes or CVD aged [Disaggregation: Age] who have had [Disaggregation: Tests] recorded within the previous 12 months.

Denominator

The number of Indigenous regular [Disaggregation: Sex at Birth] clients with Type 2 Diabetes or CVD aged [Disaggregation: Age] (see [PI05](#)).

Disaggregation

A visual representation of the disaggregation per calculation (Type 2 Diabetes, CVD and Type 2 Diabetes or CVD) can be seen below:

		Age (years)					
		18 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
eGFR only	Male	✓	✓	✓	✓	✓	✓
	Female	✓	✓	✓	✓	✓	✓
ACR only	Male	✓	✓	✓	✓	✓	✓
	Female	✓	✓	✓	✓	✓	✓
eGFR and ACR	Male	✓	✓	✓	✓	✓	✓
	Female	✓	✓	✓	✓	✓	✓
Neither an eGFR nor an ACR	Male	✓	✓	✓	✓	✓	✓
	Female	✓	✓	✓	✓	✓	✓


Notes

- 'Type 2 diabetes' – specifically excludes Type I diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, Impaired fasting glucose, and impaired glucose tolerance CVD.
- 'ACR' – refers to an albumin/creatinine ratio, or other micro albumin urine test result.
- 'An eGFR only' – a recorded eGFR result, where an ACR result HAS NOT been recorded.
- 'An ACR only' – a recorded ACR result, where an eGFR result HAS NOT been recorded.
- 'Both an eGFR and an ACR' – a recorded eGFR result AND a recorded ACR result.
- 'Neither an eGFR NOR an ACR' – an eGFR result AND an ACR result HAS NOT been recorded.


Example - Record Kidney Function Test (eGFR result) Manually

1. Open a patient's record.

2. Select **Tools > Tool Box > Renal Function Calculator**.



3. The **Renal Function Calculator** window appears.



The screenshot shows the 'Renal Function Calculator (eGFR - Abbrev. MDRD)' window. The window includes fields for Date (15/03/2019), Time (7:59:08 AM), Gender (Female), Age (51), Height, and Patient ID (31). It has tabs for Blood Glucose, Blood Pressure, CV Risk, ECG, INR, Renal Function, Respiratory, and Weight. The 'Renal Function' tab is active, showing 'eGFR (Abbreviated MDRD formula)' and 'GFR (Cockcroft-Gault formula)'. The 'Current Measurements' section has radio buttons for 'Recorded weight: 58.3 Kg As at: 15/03/2019' and 'Ideal weight:'. There are input fields for 'Se. Creatinine:' and 'eGFR:'. A 'Data' button and a 'Graph' button are also present. At the bottom, there are buttons for 'Print', 'Reference', 'Edit', 'View', 'Save', and 'Close'.

4. Enter a value for creatine to generate eGFR value and click **Save**.

Example - Add eGFR/ACR Result from An Existing hl7 File

Please refer to this Online Help Guide: [The Holding File](#).

Check

For each age and sex at birth group:

- You cannot have more regular patients with type 2 diabetes who've had a test (eGFR, ACR or both) than the total number of regular patients with type 2 diabetes.
- You cannot have more regular patients with CVD who've had a test (eGFR, ACR or both) than the total number of regular patients with CVD.
- The total number of regular patients with type 2 diabetes should be the same as the total number of regular patients in the relevant age and sex at birth groups in these nKPIs: PI05, PI07, PI08, PI15 and PI23.
- You cannot have more regular patients with CVD or type 2 diabetes than the total number of regular patients in these nKPIs: PI03 (adults), PI09 and PI16.

PI19 – Kidney Function Test Result (Type 2 Diabetes or CVD Patients)

Description

Proportion of clients with:

- Type 2 Diabetes, OR
- CVD.
- Type 2 Diabetes and/or CVD.

Whose estimated glomerular filtration rate (eGFR) AND albumin/creatinine ratio (ACR) recorded within the previous 12 months were categorised as:

- Normal risk.
- Low risk.
- Moderate risk.
- High risk.

Note: Some risk groups have more than one result classification (see description of risk categories table below).

Description of risk categories:

- Normal risk:
 - $\text{eGFR} \geq 60 \text{ mL/min/1.73 m}^2$ AND $\text{ACR} < 3.0 \text{ mg/mmol}$.
- Low risk:
 - $(\text{eGFR} \geq 45 \text{ mL/min/1.73m}^2 \text{ and } < 60 \text{ mL/min/1.73m}^2) \text{ AND } (\text{ACR} < 3.0 \text{ mg/mmol}) \text{ OR}$
 - $(\text{eGFR} \geq 60 \text{ mL/min/1.73m}^2 \text{ AND } \text{ACR} \geq 3.0 \text{ mg/mmol and } \leq 30 \text{ mg/mmol})$.
- Moderate risk:
 - $(\text{eGFR} \geq 45 \text{ mL/min/1.73m}^2 \text{ and } < 60 \text{ mL/min/1.73m}^2 \text{ AND } (\text{ACR} \geq 3.0 \text{ mg/mmol and } \leq 30 \text{ mg/mmol}) \text{ OR}$
 - $(\text{eGFR} \geq 30 \text{ mL/min/1.73m}^2 \text{ and } < 45 \text{ mL/min/1.73m}^2) \text{ AND } (\text{ACR} \leq 30 \text{ mg/mmol})$.
- High risk:
 - $(\text{eGFR} \geq 30 \text{ mL/min/1.73m}^2 \text{ AND } (\text{ACR} > 30 \text{ mg/mmol}) \text{ OR}$
 - $(\text{eGFR} < 30 \text{ mL/min/1.73m} \text{ AND any ACR result})$.

Business Rules

- eGFR results recorded in patient file defined per PI18.
- ACR results recorded in patient file defined per PI18.
- Type II Diabetes recorded in patient file defined per PI05.
- CVD recorded in patient file defined per PI18.

Calculation 1: Male

Numerator

The number of male indigenous regular clients with [Disaggregation: Chronic disease] aged [Disaggregation: Age] whose most recent eGFR and ACR results, recorded within the previous 12 months, were [Disaggregation: Results].

Denominator

The number of male indigenous regular clients with [Disaggregation: Chronic disease] aged [Disaggregation: Age] who had both an eGFR and ACR result recorded within the previous 12 months.

Calculation 2: Female

Numerator

The number of female indigenous regular clients with [Disaggregation: Chronic disease] aged [Disaggregation: Age] whose most recent eGFR and ACR results, recorded within the previous 12 months, were [Disaggregation: Results].

Denominator

The number of female indigenous regular clients with [Disaggregation: Chronic disease] aged [Disaggregation: Age] who had both an eGFR and ACR result recorded within the previous 12 months.

Disaggregation

A visual representation of the disaggregation per calculation (male and female) can be seen below:

		Age (years)					
		18 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Type 2 Diabetes	Pathology result categories (see table below)	✓	✓	✓	✓	✓	✓
		✓	✓	✓	✓	✓	✓
CVD	Pathology result categories	✓	✓	✓	✓	✓	✓
		✓	✓	✓	✓	✓	✓
Type 2 Diabetes OR CVD	Pathology result categories	✓	✓	✓	✓	✓	✓
		✓	✓	✓	✓	✓	✓

Pathology result categories

Renal function test result	Range	Applicable sex at birth
Normal Risk	eGFR $\geq$ 60 and ACR $<$ 3.0	Male
Normal Risk	eGFR $\geq$ 60 and ACR $<$ 3.0	Female
Low Risk	(eGFR $\geq$ 45 and $<$ 60 and ACR $<$ 3.0) OR (eGFR $\geq$ 60 and ACR $\geq$ 3.0 and $\leq$ 30)	Male
Low Risk	(eGFR $\geq$ 45 and $<$ 60 and ACR $<$ 3.0) OR (eGFR $\geq$ 60 and ACR $\geq$ 3.0 and $\leq$ 30)	Female
Moderate Risk	(eGFR $\geq$ 45 and $<$ 60 and ACR $\geq$ 3.0 and $\leq$ 30) OR (eGFR $\geq$ 30 and $<$ 45 and ACR $\leq$ 30)	Male
Moderate Risk	(eGFR $\geq$ 45 and $<$ 60 and ACR $\geq$ 3.0 and $\leq$ 30) OR (eGFR $\geq$ 30 and $<$ 45 and ACR $\leq$ 30)	Female
High Risk	(eGFR $\geq$ 30 and ACR $>$ 30) OR (eGFR $<$ 30 and any ACR result)	Male
High Risk	(eGFR $\geq$ 30 and ACR $>$ 30) OR (eGFR $<$ 30 and any ACR result)	Female

Notes

- 'Type 2 diabetes' – specifically excludes Type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, Impaired fasting glucose, and impaired glucose tolerance.
- The report only considers the most recently recorded eGFR test result per patient.
- The report only considers the most recently recorded ACR test result per patient.
- Results taken from relevant pathology results. If your organisation does not have a system for adding pathology results to patient records, ensure the results have been included in the correct fields for this nKPI.
- For correct data entry please see PI05.
- 'ACR' – refers to an albumin/creatinine ratio, or other micro albumin urine test result.

Example - Add eGFR data

eGFR results may be added manually through the Renal Function Calculator or via HL7 file as described in PI18.

Example - Add history of Type 2 Diabetes or Cardiovascular Disease

Type 2 Diabetes can be recorded via the method shown in PI05.

Check

For each age and sex at birth group:

- You cannot have more regular patients with type 2 diabetes with eGFR results in each range than the total number of regular patients with type 2 diabetes who had an eGFR result recorded.
- You cannot have more regular patients with CVD with eGFR results in each range than the total number of regular patients with CVD who had an eGFR result recorded.

- The total number of regular patients with type 2 diabetes who had an eGFR recorded should be the same as the sum of the number of regular patients with type 2 diabetes who had an eGFR only recorded and both eGFR and ACR recorded in this nKPI: PI18.
- The total number of regular patients with CVD who had an eGFR recorded should be the same as the number of regular patients with CVD who had an eGFR recorded in this nKPI: PI18.
- You cannot have more regular male patients with type 2 diabetes with ACR results in each specified ranged than the total number of regular male patients with type 2 diabetes who had an ACR result recorded.
- The total number of regular male patients with type 2 diabetes with an ACR result recorded should be the same as the sum of the number of regular male patients with type 2 diabetes who had an ACR only recorded and both eGFR and ACR recorded in this nKPI: PI18.
- You cannot have more regular female patients with type 2 diabetes with ACR results in each specified ranged than the total number of regular female patients with type 2 diabetes who had an ACR result recorded.
- The total number of regular female patients with type 2 diabetes with an ACR result recorded should be the same as the sum of the number of regular female patients with type 2 diabetes who had an ACR only recorded and both eGFR and ACR recorded in this nKPI: PI18.

PI20 – Cardiovascular Disease (CVD) Risk Assessment

Description

Proportion of Indigenous regular patients with no known cardiovascular disease (CVD), aged 35 to 74, with information available to calculate their absolute CVD risk.

Business Rules

- CVD recorded per PI18.
- Smoking status recorded per PI09.
- Diabetes recorded per PI05.
- Absolute CV risk is defined as patients with a measurement type of 'ACVRISK'. These results are collated into low, medium, and high by the below calculations:
 - $0 \leq \text{measurement value} < 10$: set element value as 3 (low).
 - $10 \leq \text{measurement value} \leq 15$: set element value as 2 (medium).
 - $15 < \text{measurement value}$: set element value as 1 (high).
- High-Density Lipoprotein Cholesterol recorded is defined as patients with a measurement type of 'LIPIDDATE' or 'HDL'.

Numerator

Number of regular Indigenous patients whose:

- Smoking Status has been recorded, AND
- Diabetes status, BP, Total cholesterol, and High-Density Lipoprotein Cholesterol have been measured and recorded within the last 24 months.

Denominator

Number of regular Indigenous patients **NOT CODED** with CVD (As we want Check for risk factors after) (See PI07).

Records Excluded

- Patients coded with CVD.
- Patients without known CVD if information is not available for all risk factors (tobacco smoking, diabetes, systolic blood pressure, total cholesterol and HDL cholesterol levels, age, sex).

Reporting by Age Group

Separate the results for males and females into each of the following age groups:

	Age (years)			
	35 to <45	45 to <55	55 to <65	65 to <75
Male	✓	✓	✓	✓
Female	✓	✓	✓	✓


Notes

Ensure information on the diabetes status from the most recent record is included regardless of how old that record is.


Where a patient's tobacco smoking status and/or sex does not have an assessment date assigned within the Patient Information Record System (PIRS), smoking status and/or sex as recorded in the PIRS should be treated as current (that is, as having been updated within the previous 24 months).

Example - Record Blood Pressure

1. Open a patient's record.
2. Select **Tools > Tool Box > Blood Pressure**




- The Blood Pressure module appears.




- Enter values for blood pressure and then click **Save**.

Example - Record Smoking Status

- Open a patient's record.
- Select **Patient > Details**.



3. Select the **Smoking** tab.




The screenshot shows the 'Patient Details' window with the 'Smoking' tab selected. The form contains the following fields and options:

- Date of assessment:** 14/03/2019
- Smoker:** Never smoked (dropdown)
- Frequency:** (dropdown)
- Number of cigarettes:** 5
- Year commenced:** 1980
- Duration:** 39yrs
- Stage of change assessment:** (dropdown)
- Last quit attempt:** 14/03/2019 (calendar icon) ☒ Never/Unknown
- Duration of longest period of abstinence:** (dropdown)
- ☐ Smoking cessation intervention discussed with patient
- Buttons:** View Patient Education Leaflet, Reference
- Footer:** ☐ Update address for all family members, ☐ Auto-capitalise names, Save, Cancel


4. Select the smoking status from the **Smoking** drop-down list and click **Save**.

Example - Record Total Cholesterol and HDL

1. Open a patient's record.
2. Select Clinical > Diabetes Record.



- The **Diabetes Follow Up** window appears.



Diabetes Follow Up

Assessments performed:

Assessment date: 11/11/2016

Last review by:

Ophthalmologist: 11/11/2016

Podiatrist: 11/11/2016

Dietitian: 11/11/2016

Endocrinologist: 11/11/2016

Diabetes educator: 11/11/2016


Last provided HypoKit: 11/11/2016

Buttons: Assessment, Add Values, Graph, Close

Major parameters:

Date	f Blood Gluc.	Height (cm)	Weight (kg)	HbA1c mmol/mol (%)	Creatinine
22/03/2007			69		
10/05/2007			72.0		
30/08/2007		175	65		
25/08/2008		175	63		
25/08/2010		175	66		
17/08/2012		175	64		60
11/11/2016		175	64		

- Click **Add Value**. The **Diabetes Record** window appears.



Diabetes record

Date: 14/03/2019

Parameters:

Weight (kg): Height (cm):

Systolic BP: Diastolic BP:

f Blood Gluc.: HbA1c (mmol/mol):

Total Chol.: HDL:

Triglycerides: Creatinine:

Urinalysis:

Protein:

Blood:

Glucose:

Ketones:

Leucocytes:

Bilirubin:

Last review by:

Ophthalmologist: 14/03/2019

Podiatrist: 14/03/2019

Dietitian: 14/03/2019

Endocrinologist: 14/03/2019

Diabetes educator: 14/03/2019

Last Provided HypoKit: 14/03/2019

Buttons: Save, Close

- Enter the value for total cholesterol and HDL, then click **Save**.

Check

For each age and sex at birth group:

- You cannot have more regular patients with all the following information recorded (tobacco smoking, diabetes, systolic blood pressure, total cholesterol and HDL cholesterol levels, age, sex), than the total number of regular patients without known CVD recorded.
- You cannot have more regular patients without known CVD than the total number of regular patients.

- The total number of regular patients without known CVD should be equal to the total number of regular patients in the relevant age and sex at birth groups in these nKPIs: PI03, PI09 and PI16, minus the total number of regular patients with CVD in the relevant age and sex at birth groups in this nKPI: PI18.

PI21 – Cardiovascular Disease (CVD) Risk Assessment Results

Description

Proportion of Indigenous regular patients with no known cardiovascular disease (CVD), aged 35–74, who have had an absolute CVD risk assessment recorded within the previous 24 months and whose CVD risk was categorised as one of the following:

Category	% Chance of CV Event in The Next 5 Years
Low	Less than 10% chance of a cardiovascular event in the next 5 years
Moderate	10%–15% chance of a cardiovascular event in the next 5 years
High	Greater than 15% chance of a cardiovascular event in the next 5 years

Business Rules

- CVD recorded in patient file per PI18.
- Absolute CVD risk assessment recorded per PI20.

Calculation 1: Patients with High CV Risk

Numerator:

The number of regular Indigenous patients who have had an absolute CVD risk assessment recorded in the 24 months up to the census date with risk assessed as high (greater than 15% over the next 5 years).

Denominator

Number of regular Indigenous patients identified in the Denominator include:

- Patients NOT CODED with CVD (See PI07) AND
- Having Absolute CVD risk measured within the last 24 months AND
- Aged 35 – 74.

Calculation 2: Patients with Moderate CV Risk

Numerator:

The number of regular Indigenous patients that have had an absolute CVD risk assessment recorded in the 24 months up to the census date with risk assessed as moderate (10%–15% over the next 5 years).

Denominator

Number of regular Indigenous patients identified in the Denominator include:

- Patients **NOT** CODED with CVD (See PI07) AND
- Having Absolute CVD risk measured within the last 24 months AND
- Aged 35 – 74.

Calculation 3: Patients with Low CV Risk

Numerator:

The number of regular Indigenous patients that have had an absolute CVD risk assessment recorded in the 24 months up to the census date with risk assessed as low (less than 10% over the next 5 years).

Denominator:

Number of regular Indigenous patients identified in the Denominator include:

- Patients NOT CODED with CVD (See PI07) AND
- Having Absolute CVD risk measured within the last 24 months AND
- Aged 35 – 74.

Reporting by Age Group:

Separate the results for males and females into each of the following age groups:

	Age (years)			
	35 to <45	45 to <55	55 to <65	65 to <75
Male	✓	✓	✓	✓
Female	✓	✓	✓	✓

Records Excluded


- Patients coded with CVD.
- Patients aged under 35.
- Patients aged 75 or over.

Notes


Only the most recently recorded result from an absolute CVD risk assessment. This means that if a patient has had several assessments, then include only the results from the most recent test.

Examples of Recording Risk Factors

1. From within the Clinical Window, select **Clinical > Diabetes Record**. The **Diabetes Follow Up** window appears.



2. An assessment may be conducted, or alternatively values may be added directly.



3. From within the Clinical Window, select **Tools > Toolbox > Blood Pressure**. The Blood Pressure module appears. Record Blood Pressure from within this window.

Tool Box

✕

Blood Pressure

Date:
Time:
Gender:
Age:
Height:
Patient ID: 31

Blood Glucose

Blood Pressure

CV Risk

ECG

INR

Renal Function

Respiratory

Weight

Device

Manual

Data

Graph

Current Measurements

View:

Blood Pressure

Cuff Location:

Systolic / Diastolic

Pulse

Sitting:

Standing:

Lying:

Date	Time	Location	Type	BP	Pulse	Rhythm

<

>

Clear

Record

Print

Reference

Edit

View

Save

Close

Check

For each age and sex at birth group:

- You cannot have more regular patients without known CVD with absolute CVD risk assessment results in each range than the total number of regular patients without known CVD who had an absolute risk assessment recorded.
- You cannot have more regular patients without known CVD than the total number of regular patients.
- The total number of regular patients without known CVD who had an absolute risk assessment recorded should be less than or equal to the number of patients without known CVD who had all the following information (tobacco smoking, diabetes, systolic blood pressure, total cholesterol and HDL cholesterol levels, age, sex) recorded in this nKPI: PI20.

PI22 – Cervical Screening Recorded

Description

Proportion of female regular patients who are Indigenous, aged 25–74, who have not had a hysterectomy and who have had a cervical screening (human papillomavirus (HPV)) test within the previous 5 years.

This indicator has been revised to align with the new National Cervical Screening Program (NCSP) where the previous Pap test is replaced by an HPV test from 1 December 2017. For the nKPIs these changes take effect from June 2018 reporting period.

The key changes are:

- Data is to be collected on patients who had a human papillomavirus (HPV test) conducted from 1 December 2017.
- The HPV test can be based on a sample collected by a health practitioner or on a self-collected sample.
- The age range for this nKPI indicator has been revised to 25–74 for a transitional period.
- Data is to be disaggregated by age group (see specifications below).

Business Rules

- Cervical screening as recorded a record present in the 'Cervical Screening' tab.
- No history of hysterectomy is defined as patients whose medical file does not contain the below DOCLE codes:

cxyz	Hysterectomy
cxyz	Surgery - Uterus - Hysterectomy
cxyz	Uterus - removal of
cxyz	Hysterectomy - Abdominal
cxyz	Hysterectomy - Vaginal
cxyz	Vaginal Hysterectomy
cxyz	Hysterectomy - Vaginal with vaginal repair
cxyz	Hysterectomy - Laparoscopic
cxyz	Laparoscopic hysterectomy
cxyz	Hysterectomy - ovary(ies) spared
cxyz	Hysterectomy - Subtotal
cxyz	Hysterectomy - Total
cxyz	Subtotal hysterectomy
cxyz	Hysterectomy & BSO - Abdominal

Numerator

Number of female Indigenous patients with cervical screening done within the previous 5 years.

Denominator

Number of regular Indigenous female patients with no history of hysterectomy aged 25 – 74.

Records Excluded

- Patients whose last screening was more than 5 years ago.
- Patients with hysterectomy present in their record as a history item or reason for visit. See above for hysterectomy DOCLE Codes.

Reporting by Age Group

Results should be separated into the below age groups:


	Age (years)				
	25 to <35	35 to <45	45 to <55	55 to <65	65 to <75
Female	✓	✓	✓	✓	✓

Notes

Include patients who had HPV tests conducted from 1 December 2017, HPV tests where the sample is either collected by a health practitioner or self-collected and results from other organisations (where possible).

Example – Adding a Cervical Screening Result

1. Open a female patient's record.
2. Select the **Cervical Screening** tab.
3. Right-click in the white space and select **New Item**.



4. The **Record Cervical Screening Result** window appears. Enter the date of the result, select the result from the drop down and tick the 'Endocervical cells present?' and/or HPV changes present?' checkboxes as required.

Record Cervical Screening Result

Jennifer ANDREWS
DOB: 20/04/1967 Record No:

Screening Result

Date: 15/03/2019 Result: Negative

☐ Endocervical cells present? ☐ HPV changes present?

Comment:

[View AMBS 2004 Comparison Table](#)

5. Click **Add**.

Check

- The number of female regular patients aged 25–74 with a cervical screening should not exceed the total number of female regular patients aged 25–74 who have not had a hysterectomy.

PI23 – Blood Pressure Recorded (Type 2 Diabetes Patients)

Description

Proportion of regular patients who are Indigenous, have type 2 diabetes and who have had a blood pressure measurement result recorded at the primary health-care organisation within the previous 6 months.

Business Rules

- Type 2 Diabetes recorded as per PI05.
- Blood pressure measurement recorded as per patient's whose file's contain measurements of types 'Systolic' and 'diastolic' as recorded in the 'Blood pressure' toolbox.

Numerator

Number of Indigenous patients with BP recorded within the past 6 months, separated into the below age groups (See PI20).

Denominator

Number of Indigenous patients with Diabetes Type 2 (See PI05).

Records Excluded


Patients with type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose or impaired glucose tolerance.

Reporting by Age Group


	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Example - Record Blood Pressure

1. Open a patient's record.
2. Select **Tools > Tool Box > Blood Pressure**.



3. The **Blood Pressure** module appears.



4. Enter values for Systolic and Diastolic and click **Save**.

Check

For each age and sex at birth group with type 2 diabetes:

- You cannot have more regular patients with type 2 diabetes who've had a blood pressure test in the past 6 months than the total number of regular patients with type 2 diabetes.
- The number with type 2 diabetes who've had a blood pressure test should be the same as the total number of regular patients with type 2 diabetes who had a blood pressure test in this nKPI: PI24.
- The total number of regular patients with type 2 diabetes should be the same as the total number of regular patients with type 2 diabetes in the relevant age and sex at birth groups in these nKPIs: PI05, PI07, PI08, PI15 and PI18.
- The total number of regular patients with type 2 diabetes cannot be more than the total number of regular patients in these nKPIs: PI03 (adults), PI09 and PI16.

PI24 – Blood pressure less than or equal to 140/90 mmHg (type 2 diabetes patients)

Description

Proportion of regular patients who are Indigenous, have type 2 diabetes and whose blood pressure measurement result, recorded within the previous 6 months, was less than or equal to 140/90 mmHg.

Business Rules

- Type 2 Diabetes as recorded in PI05.
- Blood pressure recorded per PI23.

Numerator

Number of male and female indigenous patient records in each age group below, with a recorded blood pressure of 140/90mmHg or less in the 6 months up to the census date.

Denominator

Number of indigenous patient records identified in the Denominator include:

- Patients with Diabetes Type 2 (See PI05) AND
- Has BP recorded within the past 6 months? (See PI20).

Reporting by Age Group

	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Records Excluded

Any patients whose systolic or diastolic reading is above the threshold (140 and 90, respectively) in the Numerator. (They will be counted in the total number who have had a blood pressure test.)

- Patients with type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose or impaired glucose tolerance.

Notes

- Only the most recently recorded blood pressure test result. This means that if a patient has had their blood pressure measured several times in the past 6 months, then include only the results from the most recent measurement.

Check

For each age and sex at birth group of regular patients with type 2 diabetes:

- The number whose blood pressure is 140/90mmHg or less cannot be more than the number who've had their blood pressure tested.
- The total number with type 2 diabetes who had a blood pressure test should be the same as the number of regular patients with type 2 diabetes who had a blood pressure test in this nKPI: PI23

PI25 – STI Result Recorded

Description

Proportion of Indigenous regular clients aged 15 to 34 years who had a STI(s) result recorded (chlamydia and/or gonorrhoea) within the previous 12 months.

Business Rules

Patients with either a Chlamydia or Gonorrhoea test result recorded is defined as patients whose records contain a pathology result with a relevant LOINC codes (Appendix B) or Wild card search for a valid test name (Appendix C).

Numerator

Number of male and female indigenous patient records in each age group below, who have had a chlamydia and/or gonorrhoea test result recorded within the previous 12 months.

Denominator

Number of male and female indigenous patient records in each age group below.

Reporting by Age Group

	Age (years)			
	15 to <20	20 to <25	25 to <30	30 to <35
Male	✓	✓	✓	✓
Female	✓	✓	✓	✓

Check

For each age and sex at birth group of regular patients with an STI result recorded:

- The number of tests recorded in each cell cannot be more than the number of patients in each age group.

PI26 – Ear Health Check

Description

Proportion of Indigenous regular clients who, in the previous 12 months, have had a completed check, that was recorded and classified into:

- Appearance of both ear canals and eardrums
- Movement of both eardrums
- Appearance of both ear canals and ear drums AND movement of both* ear drums

Business Rules

- Clients may be counted in one or all parts of the indicator.
- A recorded ear health check in the last 12 months is determined from the following workflow, in the following order:
 - i. Ear check recorded in the ear health section or 'module' in the CIS – if recorded here do not count 'ii' and 'iii'
 - ii. Search for procedure codes and terms as defined in the ear health coding framework (Appendix A)
 - iii. If none are found, search for condition codes and terms as defined in the ear health coding framework (Appendix A)
- Count only conditions and procedures recorded in the last 12 months.
- For workflow 'ii' and 'iii', count procedures and/or conditions against either Appearance, Movement or BOTH Appearance and Movement as defined in the ear health coding framework.
- Currently, MDInsights supports workflow ii and iii.

Calculation 1: Patients with Appearance of both ear canals and eardrums

Numerator:

Number of *[Disaggregation: Sex at Birth]* Indigenous regular clients aged *[Disaggregation: Age]* who have had a completed check of the **appearance of both ear canals and ear drums** within the previous 12 months.

Denominator

Total number of *[Disaggregation: Sex at Birth]* Indigenous regular clients aged *[Disaggregation: Age]*.

Calculation 2: Patients with Movement of both eardrums

Numerator:

Number of *[Disaggregation: Sex at Birth]* Indigenous regular clients aged *[Disaggregation: Age]* who have had a completed check of the **movement of both eardrums** within the previous 12 months.

Denominator

Total number of *[Disaggregation: Sex at Birth]* Indigenous regular clients aged *[Disaggregation: Age]*.

Calculation 3: Patients with Appearance of both ear canals and ear drums AND movement of both ear drums

Numerator:

Number of *[Disaggregation: Sex at Birth]* Indigenous regular clients aged *[Disaggregation: Age]* who have had a completed check of the **appearance of both ear canals and eardrums and the movement of both eardrums** within the previous 12 months.

Denominator

Total number of *[Disaggregation: Sex at Birth]* Indigenous regular clients aged *[Disaggregation: Age]*.

Reporting by Age Group

	Age					
	0 to <12 Months	12 to <24 Months	24 to <36 Months	36 to <60 Months	5 to <10 Years	10 to <15 Years
Male	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓

Check

- For each age and sex at birth group of regular patients with an ear health check recorded should not be more than the total number of regular clients in that group.

APPENDIX A- DOCLE CODES

Diabetes Type 2

diabm@niddm	Diabetes Mellitus – NIDDM
diabm@niddm	Diabetes Mellitus - Type II
diabm@niddm	NIDDM
diabm@niddm	NIDDM (Non-Insulin dependent diabetes mellitus)
diabm@niddm	Non-insulin dependent diabetes mellitus
diabm@niddm&rx%insu	Diabetes Type II requiring insulin
diabm@niddm&rx%insu	NIDDM - requiring insulin

Unspecified Diabetes

diabm	Diabetes Mellitus
-------	-------------------

COPD

bronchit@chro	Bronchitis – Chronic
coad	CAL (Chronic Airways Limitation)
coad	Chronic Airways Limitation
coad	Chronic Bronchitis
coad	Chronic Obstructive Airways Disease
coad	Chronic Obstructive Pulmonary Disease
coad	COAD
coad	COAD (Chronic Obstructive Airways Disease)
coad	COPD (Chronic Obstructive Pulmonary Disease)
emph	Emphysema

CVD

Coronary Heart Disease (CHD)

Atherosc	Arteriosclerosis
Atherosc	Atherosclerosis
embo.arte	Arterial embolism
embo.arte.aort	Aortic embolism
Ihd	Atherosclerotic heart disease
Ihd	Coronary artery disease
Ihd	Coronary heart disease
Ihd	Coronary insufficiency
Ihd	IHD
Ihd	IHD (Ischaemic Heart Disease)

Coronary Heart Disease (CHD)

lhd	Ischaemic heart disease
lhd	Myocardial insufficiency
occl@coroa	Blockage Coronary Artery
occl@coroa	Coronary Occlusion
occl@coroa	Occlusion-Coronary Artery
ppoc@reha@ihd	Coronary artery disease-Rehabilitation

Acute Coronary Syndrome (ACS)

Angip	Angina
Angip	Angina pectoris
angip@unst	Acute coronary insufficiency
angip@unst	Angina pectoris – unstable
angip@unst	Preinfarction syndrome
angip@unst	Unstable Angina
angip@unst:high@risk	Unstable Angina - High risk
angip@unst:low@risk	Unstable Angina - Low risk
angip@unst:moderate@risk	Unstable Angina - Moderate risk
occl@coroa:acut	ACS (acute coronary syndrome)
occl@coroa:acut	Acute coronary syndrome
occl@coroa:acut	Acute Coronary Syndrome (ACS)

Carotid Artery Stenosis

sten.caroa	Carotid Stenosis
------------	------------------

Surgery – Cardiovascular – Carotid

surg.caroa@bypa@graf	Carotid artery bypass surgery
surg.caroa@enda	Carotid Endarterectomy
surg.caroa@enda	Endarterectomy – Carotid
surg.caroa@stent	Arterial stent - Carotid artery
surg.caroa@stent	Carotid artery stent
surg.caroa@stent	Stent - Carotid artery

Surgery – Cardiovascular – Coronary

surg.coroa	Coronary artery surgery
surg.coroa	Surgery - Coronary artery
surg.coroa@balloon	Angioplasty – coronary
surg.coroa@balloon	Balloon coronary angioplasty
surg.coroa@balloon	Coronary artery balloon angioplasty
surg.coroa@balloon	PCTA
surg.coroa@balloon	Percutaneous transluminal angioplasty
surg.coroa@balloon	Surgery - Coronary artery balloon angioplasty
surg.coroa@enda	Coronary artery endarterectomy

surg.coroa@enda	Surgery - Coronary artery endarterectomy
surg.coroa@graf	Bypass – coronary
surg.coroa@graf	CABG (Coronary Artery Bypass graft)
surg.coroa@graf	Coronary artery bypass graft
surg.coroa@graf	Surgery - Coronary artery bypass graft
surg.coroa@stent	Coronary Angiography and Stent
surg.coroa@stent	Coronary artery stent
surg.coroa@stent	Stent-Coronary artery
surg.coroa@stent	Surgery - Coronary artery stent
surg.coroa@stent	Angioplasty - coronary (with stent)
surg.coroa@stent@eluting	Arterial stent - Coronary artery, drug-eluting
surg.coroa@stent@eluting	Coronary artery, drug-eluting stent
surg.coroa@stent@eluting	Stent-Coronary Artery, drug-eluting
surg.coroa@stent@eluting@non	Arterial stent - Coronary artery, not drug-eluting
surg.coroa@stent@eluting@non	Coronary artery ,not drug-eluting stent
surg.coroa@stent@eluting@non	Stent-Coronary artery, not drug-eluting

Cerebrovascular Disease

Cva	Cerebral Infarction
Cva	Cerebrovascular Accident
Cva	CVA (Cerebrovascular Accident)
Cva	Stroke
cva@isch	Ischaemic Stroke
cva@isch	Stroke – Ischaemic
cva@lacu	Lacunar Stroke
cva@lacu	Stroke – Lacunar
cva@migr	Migranous Stroke
cva@migr	Stroke – Migranous
cva@thrombos	Stroke – Thrombotic
cva@thrombos	Thrombotic – Stroke
Cvi	Cerebrovascular insufficiency
embo.cere	Cerebral Embolism
embo.cere	Embolism – Cerebral
Ich	Cerebral Haemorrhage
Ich	Haemorrhage – intracerebral
Ich	Haemorrhagic CVA
Ich	Haemorrhagic Stroke
Ich	Intracerebral bleed
Ich	Intracerebral haemorrhage
Ich	Stroke-Haemorrhagic
Sah	Haemorrhage – subarachnoid
Sah	Subarachnoid bleed
Sah	Subarachnoid haemorrhage

Cerebrovascular Disease - TIA

Tia	Arterial Embolism – minor
Tia	Cerebral TIA
Tia	Cerebral Transient Ischaemia
Tia	Cerebral Transient Ischaemic Attacks
Tia	TIA (Transient Ischaemic Attack)
Tia	Transient Ischaemic Attack

Myocardial Infarction (MI)

Myoci	Acute myocardial infarction
Myoci	AMI (Acute Myocardial Infarction)
Myoci	Myocardial Damage
Myoci	Myocardial Damage
Myoci	Myocardial infarction
Myoci	Myocardial Infarction(MI)
myoci:sans,st@elev	Myocardial infarction-without ST elevation
myoci:sans,st@elev	Non-ST-elevation myocardial infarction (NSTEMI)
myoci:sans,st@elev	NSTEMI (non-ST-elevation myocardial infarction)
myoci:st@elev	Myocardial Infarction-with ST elevation
myoci:st@elev	STEMI (ST-Elevation Myocardial Infarction)
myoci@ante	Anterior myocardial infarct
myoci@ante	Myocardial infarction-anterior
myoci@antel	Anterolateral myocardial infarct
myoci@antel	Myocardial infarction-anterolateral
myoci@antel	Myocardial infarction-anterolateral
myoci@inf	Inferior myocardial infarction
myoci@inf	Inferior myocardial infarction
myoci@inf	Myocardial infarction-inferior
myoci@inf	Myocardial infarction-inferior
myoci@posterio	Myocardial infarction-posterior
myoci@posterio	Myocardial infarction-posterior
myoci@posterio	Posterior myocardial infarct
myoci@posterio	Posterior myocardial infarct
myoci@sile	Myocardial infarction-silent
myoci@sile	Myocardial infarction-silent
myoci@sile	Silent myocardial infarction
myoci@sile	Silent myocardial infarction
myoci@subendoc	Myocardial infarction-subendocardial
myoci@subendoc	Myocardial infarction-subendocardial
myoci@subendoc	Subendocardial myocardial infarct
myoci@supe	Myocardial infarction-superior
myoci@supe	Myocardial infarction-superior
myoci@supe	Superior myocardial infarct

Peripheral Vascular Disease (PVD)

Pvd	Arteriosclerosis obliterans
Pvd	Peripheral Vascular Disease
Pvd	PVD

Renal Artery Stenosis (RENALSTENOSIS)

sten.renaa	Renal Artery Stenosis
sten.renaa	Stenosis - Renal artery

Surgery – Cardiovascular – Renal Artery

surg.renaa	Angioplasty – renal
surg.renaa@stent	Arterial stent - Renal artery
surg.renaa@stent	Renal artery stent
surg.renaa@stent	Stent - Renal artery

Chronic Kidney Disease

crf:eval,ctx:stag3	Chronic Kidney Disease - Stage 3
crf:eval,ctx:stag3	CKD (Chronic Kidney Disease) Stage 3
crf:eval,ctx:stag3	Kidney Disease - Chronic - Stage 3
crf:eval,ctx:stag3	Renal Disease - Chronic - Stage 3
crf:eval,ctx:stag4	Chronic Kidney Disease - Stage 4
crf:eval,ctx:stag4	CKD (Chronic Kidney Disease) Stage 4
crf:eval,ctx:stag4	Kidney Disease - Chronic - Stage 4
crf:eval,ctx:stag4	Renal Disease - Chronic - Stage 4
crf:eval,ctx:stag5	Chronic Kidney Disease - Stage 5
crf:eval,ctx:stag5	CKD (Chronic Kidney Disease) Stage 5
crf:eval,ctx:stag5	Kidney Disease - Chronic - Stage 5
crf:eval,ctx:stag5	Renal Disease - Chronic - Stage 5
ppoc@capd	CAPD (Continuous Ambulatory Peritoneal Dialysis)
ppoc@capd	Continuous Ambulatory Peritoneal Dialysis
ppoc@capd	Dialysis peritoneal
ppoc@capd	Peritoneal Dialysis
ppoc@dial@kidn	Dialysis haemodialysis
ppoc@dial@kidn	Haemodialysis
ppoc@dial@kidn	Hemodialysis
ppoc@dial@kidn	Renal dialysis
surg.kidn@transpla	Kidney transplant
surg.kidn@transpla	Renal transplant
surg.kidn@transpla	Surgery - Kidney transplant
surg.kidn@transpla	Transplant – kidney

Hysterectomy

Cxyz	Hysterectomy
Cxyz	Surgery - Uterus – Hysterectomy
Cxyz	Uterus - removal of
Cxyz	Hysterectomy – Abdominal
Cxyz	Hysterectomy – Vaginal
Cxyz	Vaginal Hysterectomy
Cxyz	Hysterectomy - Vaginal with vaginal repair
Cxyz	Hysterectomy – Laparoscopic
Cxyz	Laparoscopic hysterectomy
Cxyz	Hysterectomy - ovary(ies) spared
Cxyz	Hysterectomy – Subtotal
Cxyz	Hysterectomy – Total
Cxyz	Subtotal hysterectomy
Cxyz	Hysterectomy & BSO – Abdominal

Ear Health – Conditions

Otite	Otitis Externa	Both
otite@recu	Otitis Externa - Recurrent	Both
Otite	Outer ear infection	Both
otitm&ctx@ill,recu	Chronic Suppurative Otitis Media	Both
Otitm	Middle ear infection	Both
Otitm	Otitis Media	Both
otitm&ctx@ill,recu	Otitis Media - Recurrent	Both
tube.ear	Ear tuberculosis	Both
myringit<vuru	Myringitis	App
perf.eard	Tympanic perforation	App
perf.eard	Ear drum perforation	App
perf.eard	Perforation - eardrum	App
perf.eard	Perforation of eardrum	App
Cholstea	Cholesteatoma	App
effu.ear	Ear effusion	App
ear@disc	Ear - Discharge	App
ear@disc	Otorrhoea	App
ear@blee	Ear - Bleeding	App
ear@blee	Bleeding from ear	App

Ear Health – Procedures

surg.ear@tympm@perf	Myringoplasty	Both
surg.ear@tympm@inci	Myringotomy	Both
surg.ear@tympm	Tympanic membrane surgery	Both
surg.ear@tympm@perf	Tympanoplasty	Both
<u>surg.ear@grom</u>	Tympanostomy tube insertion	Both

surg.ear@grom@remo	Tympanostomy tube removal	Both
surg.ear@grom	Insertion of grommets	Both
surg.ear@grom	Ear - grommet insertion	Both
surg.ear@grom	Grommet insertion	Both
surg.ear@grom	Surgery - Ear - Grommet insertion	Both
surg.ear@grom	Ear Grommets	Both
surg.ear@grom	Tubes - middle ear - insertion	Both
surg.ear@grom@remo	Removal of grommets	Both
surg.ear@grom@remo	Ear - grommet removal	Both
surg.ear@grom@remo	Grommet removal	Both
surg.ear@grom@remo	Surgery - Ear - Grommet removal	Both
surg.ear@grom@remo	Tubes - middle ear - removal	Both
surg.ear@tympm@inci@suct	Suction Myringotomy	Both
surg.ear@tympm	Surgery - Ear - tympanic membrane	Both
surg.ear@tympm@perf	Surgery - Ear - Tympanic membrane - Perforation repair	Both
ppoc@exam@ear	Otoscopy	App
ppoc@exam@ear	Ear examination	Both
ppoc@exam@ear@pneumati	Pneumatic otoscopy	Move
ppoc@exam@ear@endos	Ear endoscopy	App
ppoc@exam@ear@tympanom	Tympanometry	Move

APPENDIX B- LOINC CODES

Chlamydia and gonorrhoea - Included

100356-5	100710-3	101172-5	36902-5	36903-3	43405-0	43406-8
44806-8	44807-6	45067-6	45068-4	45069-2	45070-0	45072-6
45074-2	45075-9	45076-7	47362-9	64017-7	70161-5	70162-3
70163-1	70164-9	72828-7	77577-5	80360-1	80361-9	80362-7
80365-0	85687-2	85689-8	90357-5	90358-3	92683-2	92684-0
96612-7	101696-3	103633-4	106058-1	45071-8		

Chlamydia and gonorrhoea - Included local unpublished codes

39994369	M112	M1005	M1015	M2845	1230005015	D110
----------	------	-------	-------	-------	------------	------

Chlamydia and gonorrhoea - Excluded

45073-4	62864-4	62865-1	69865-4	86661-6
---------	---------	---------	---------	---------

Chlamydia only - Included

14463-4	14464-2	14465-9	14467-5	14470-9	14471-7	14472-5
14474-1	14509-4	14510-2	14511-0	14513-6	16593-6	16600-9
16601-7	21189-6	21190-4	21191-2	21192-0	21613-5	23838-6
31765-1	31767-7	31771-9	31772-7	31775-0	31776-8	31777-6
32001-0	32004-4	32671-0	32774-2	34708-8	34709-6	34710-4
35412-6	35711-1	35713-7	35714-5	35715-2	35716-0	35717-8
35718-6	35721-0	35724-4	35725-1	35726-9	35727-7	35728-5
35729-3	35730-1	35736-8	42931-6	43304-5	43404-3	45078-3
45080-9	45084-1	45085-8	45086-6	45089-0	45090-8	45091-6
45092-4	45093-2	45095-7	45096-5	45097-3	45098-1	45100-5
45101-3	45103-9	45104-7	45105-4	45106-2	45107-0	45108-8
45112-0	45113-8	45114-6	45115-3	45131-0	45132-8	45133-6
47211-8	47212-6	49096-1	4993-2	50311-0	50387-0	50411-8
51578-3	53925-4	53926-2	557-9	558-7	559-5	560-3
561-1	57287-5	57288-3	6345-3	6346-1	6347-9	6349-5
6354-5	6355-2	6356-0	6357-8	80363-5	80364-3	80367-6
82306-2	88221-7	89648-0	91860-7	91873-0	14466-7	14473-3
14512-8	16602-5	20993-2	31773-5	32002-8	35719-4	35720-2

45079-1	45081-7	45082-5	45083-3	45087-4	45088-2	45102-1
45110-4	45111-2	6358-6	104722-4	104815-6	105339-6	105340-4

Chlamydia only - Included local unpublished codes

30677925	30677929	30677933	M1010	M1020	M2850	ChlamMethTest
CHLAMYDIA PCR URINE	CHTPCR	CLR01	CLR02	D0111	D0116	D0124
D0351	D0353	D0355	D0843	D0847	D0853	D0854
D0859	D353	D841	D843	D851	D852	D853
cP						

Gonorrhoea only - Included local unpublished codes

30678089	30678097	30678101	30678101	M1011	M1022	M2852
GonoMethTest	GONO URINE PCR	GONPCR	GOLCX	D0114	D0119	D0129
D0401	D0403	D0405	D0845	D0849	D0857	D0858
D0861	D403	D845	D846	D855	D855	D856
D857	gN					

Chlamydia only - Excluded

100120-5	100121-3	13217-5	13218-3	13219-1	13220-9	13221-7
13222-5	13223-3	13224-1	14199-4	14200-0	14201-8	14202-6
14203-4	14204-2	14461-8	14462-6	5088-0	14468-3	14469-1
5089-8	14507-8	14508-6	46176-4	16594-4	16595-1	16596-9
16597-7	16598-5	16599-3	45130-2	18490-3	18491-1	18492-9
45135-1	21187-0	21188-8	22187-9	22188-7	22189-5	22190-3
22191-1	22192-9	22193-7	22194-5	22195-2	22196-0	22197-8
22198-6	22199-4	22200-0	22201-8	22202-6	26626-2	26663-5
26664-3	26665-0	26666-8	26667-6	26668-4	26715-3	27167-6
27185-8	27368-0	27370-6	27371-4	28556-9	28557-7	28558-5
30204-2	31293-4	31294-2	31295-9	31296-7	31297-5	31298-3
31768-5	31769-3	31770-1	45094-0	31774-3	32005-1	32006-9
32007-7	33574-5	33575-2	33604-0	33605-7	38469-3	40710-6
40854-2	40855-9	40856-7	41157-9	43058-7	43059-5	43060-3
43061-1	43062-9	43173-4	43174-2	43175-9	43355-7	43356-5
43357-3	43848-1	44005-7	44079-2	44983-5	44984-3	44985-0
44986-8	44987-6	44988-4	44989-2	44990-0	44991-8	44992-6
44993-4	44994-2	44995-9	44996-7	44997-5	44998-3	44999-1

45000-7	45001-5	45002-3	45003-1	45004-9	45005-6	45006-4
45007-2	45008-0	45009-8	45077-5	5090-6	5087-2	47234-0
46177-2	46178-0	43216-1	54044-3	74241-1	35639-4	33943-2
51734-2	6350-3	6351-1	6352-9	6353-7	24402-0	6918-7
6919-5	6920-3	7824-6	87949-4	87950-2	90361-7	90362-5
90363-3	90365-8	91861-5	99105-9	99778-3	10848-0	10849-8
16589-4	16590-2	16591-0	16592-8	20755-5	20756-3	20757-1
21185-4	21186-2	22182-0	22183-8	22184-6	22185-3	22186-1
23967-3	23990-5	24005-1	24238-8	24239-6	25369-0	27273-2
29664-0	29677-2	31292-6	31763-6	31764-4	31766-9	7822-0
32003-6	33410-2	34264-2	35710-3	35712-9	82957-2	90763-4
35722-8	35723-6	38467-7	38468-5	41482-1	43063-7	43846-5
43847-3	45010-6	45011-4	45012-2	45099-9	39224-1	45109-6
96075-7	57817-9	46179-8	5082-3	5083-1	5084-9	5085-6
5086-4	556-1	56909-5	56910-3	56911-1	57679-3	57680-1
6342-0	6343-8	6344-6	6348-7	75757-5	77166-7	7823-8
86659-0	100122-1	100123-9	100124-7	10651-8	10652-6	11254-0
16581-1	16582-9	16583-7	16584-5	16585-2	21184-7	22169-7
22170-5	22171-3	22172-1	22173-9	22174-7	24004-4	29722-6
30205-9	31299-1	31300-7	31301-5	31302-3	31303-1	31304-9
31778-4	34645-2	34707-0	43844-0	43845-7	44080-0	44979-3
44980-1	44981-9	44982-7	47052-6	50612-1	5078-1	51806-8
51807-6	58758-4	6912-0	6913-8	6914-6	7821-2	82178-5
88546-7	88547-5	88718-2	89640-7	89641-5	92133-8	92577-6
92859-8	92860-6	92986-9	100125-4	14198-6	16586-0	16587-8
16588-6	20752-2	20753-0	20754-8	22175-4	22176-2	22177-0
22178-8	22179-6	22180-4	22181-2	22992-2	22993-0	22994-8
22995-5	22996-3	22997-1	22998-9	22999-7	23000-3	23001-1
27404-3	31305-6	31306-4	31307-2	31779-2	31780-0	43057-9
43868-9	44081-8	44977-7	44978-5	5079-9	5080-7	5081-5
51808-4	51809-2	59065-3	6338-8	6339-6	6340-4	6341-2
6915-3	6916-1	6917-9				

Gonorrhoea only - Included

14127-5	21414-8	21415-5	21416-3	24111-7	29311-8	31905-3
31906-1	32198-4	32199-2	32705-6	43305-2	43403-5	47387-6
5028-6	50326-8	50388-8	50412-6	53879-3	53927-0	57180-2
57289-1	57458-2	60255-7	60256-5	6487-3	6488-1	6489-9
6490-7	688-2	690-8	691-6	692-4	693-2	694-0
696-5	697-3	698-1	80366-8	80368-4	88224-1	88225-8
96599-6	104765-3	104780-2	105085-5	105922-9	106572-1	23908-7
39557-4	53878-5	6491-5	689-0			

Gonorrhoea only - Excluded

22430-3	695-7	27021-5	30099-6	31525-9	32704-9	33904-4
35735-0	86660-8	42987-8	43387-0	5261-3	53762-1	9568-7
91781-5	97626-6	99779-1				

APPENDIX C- FREE TEXT SEARCH TERMS FOR PI25

Chlamydia - Included

Chlamydial infection	Chlamydial STI	Chlamydia trachomati
Chlamydia trachomatis	Chlamydial trachomatis	c trachom
Chlamydial	Chlamydia	Chlam
ct main	CTNG	CT/NG
CHL/GON	U-CHL/GON	NG and CT
NGCT	c.trachomatis	CHPCRT

Chlamydia - Excluded

Pneumoniae	Pneum	Psittaci
Ornithosis	Orni	Ab
IgA	IgG	IgM

Gonorrhoea - Included

Gono	Gonno	Neisseria gonorrhoea
Gonococcus	Gonorrhea	N gonorrhoea
Gonorrhoea	Gonorrhoeae	Gnrrh
ng main	CTNG	CT/NG
CHL/GON	U-CHL/GON	NG and CT
NGCT	NGPCR	NGPCRT


Gonorrhoea - Excluded

Ab	IgA	IgG
IgM		

APPENDIX D- CREATING AND PROCESSING A CLAIM

Example – Creating a 715 Claim

1. Open MedicalDirector Pracsoft.
2. Add Patient to the Pracsoft **Waiting Room**.
 - See '[Adding Patients to the Waiting Room](#)' for instructions.
3. Right-click the patient's name, and click **Record Visit**



4. Select the following:
 - Select **Bulk Bill** via the **Invoice To** drop-down list.
 - Select the associated Doctor via the **Seen By** drop-down list.
 - Service Details: Item No: 715.

Record Visit

File View

Patient name: ANDREWS, Jennifer Seen by: AP Location: Surgery

Consult time: 10:18:42 AM

Consult date: 15/03/2019

Invoice to: Bulk Bill

Visit Date	Item No	Description	Fee	Gap	Fee Rate	GST Rate	GST Amount	Text
15/03/2019	715	attendance by a medical practitioner...	\$176.90	\$0.00	Rebate 85	0.00	\$0.00	

\$0.00 \$0.00 \$0.00

☐ In hospital ☐ Apply incentive

Add Delete Up Down

Payment options (F12)

Pay now Full Payment

Pay later Gap Payment

Visit options

☒ Remove from Waiting Room ☐ Non visit

☐ Non patient account ☐ Aftercare

Print options

☒ Print voucher ☒ Print preview

☐ Print PC1 Copies: 1

Hold Voucher Account Cancel

- Click **Voucher**

Example - Processing a Patient Claim

- Select **Processing > Batch Bulk Claims > Medicare**.

Patient Waiting Room Accounts Processing Reports MedicalDirector Admin Setup SMS Window Help

Banking Batch Bulk Claims Receipt Bulk Payments Medicare Veterans Affairs

All AP

Name	Chart	Appointment Time	Arrival Time	Waiting Time	Concession
ANDREWS, Jennifer			15/03/2019 10:00 am	27 mins	None

- Enter a Claim Number (5 digits).

Batch Bulk Claims - Medicare

Claim details Claim 1 of 1

Practitioner: AP Number of vouchers: 1

Location: Surgery Total amount claimed: \$176.90

Claim number: A2345

Show Process

Summary


Undaimed visits: 1

Undaimed amount: \$176.90

Minimum number of claims required: 1

Send Close

3. Click **Process**.
4. Click **Yes**.
5. Click **Send**.
6. Click **Yes** to proceed. The **Select Claims Tab** is presented.



Medicare Australia Online Claiming / Easyclaim Reconciliation

File View Setup

1 Select Claims 2 Prepare Claim Data 3 Print Claim Reports 4 Transmit Claims 5 Request Reports

Claims selected to prepare:

Claim No	Amount	Practitioner	Provider No	Type	Status	Date Lodged
A000060	\$0.00	D7F	022364BA	AIR	New	
A000064	\$0.00	JB	0385238Y	AIR	New	
A000068	\$0.00	JB	0385238Y	AIR	New	
A000072	\$0.00	JB	0385238Y	AIR	New	
A0021	\$0.00	NP1	2406901F	Medicare	New	
A0024	\$0.00	56	2406901F	Medicare	New	
A0026	\$0.00	AL	2406901F	Medicare	New	


Claims prepared/transmitted:

Claim No	Amount	Practitioner	Provider No	Type	Status	Date Lodged
#0012	\$37.05	AP	0385238Y	Medicare	Trans	1/12/2016 10:3
A0047	\$6.15	DOG	2406901F	Medicare	Trans	8/11/2016 2:41
A0054	\$37.05	AP	0385238Y	Medicare	Trans	8/11/2016 2:41
A0055	\$37.05	AP	0385238Y	Medicare	Trans	20/09/2016 10:
A0056	\$36.30	AL	2406901F	Medicare	Prep	
A0058	\$43.20	AP	2054781W	Medicare	Trans	20/09/2016 10:
A0060	\$326.92	GP	2146681K	Medicare	Prep	
A0062	\$343.15	JMT	4513946K	Medicare	Trans	8/11/2016 2:41
A0064	\$620.10	AP	0385238Y	Medicare	Trans	10/11/2016 4:1

Location Id: H5588888

Next Step Close

7. Click **Next Step**. The **Prepare Claim Tab** is presented. Click **Prepare**.
8. When prompted, click **Yes** for claim item.



PracSoft - Prepare Claims

You have selected to prepare the following Claims for the following Practitioners

A5222


AP (2426621B)

Do you wish to continue?

Yes No

9. Click **Next Step**. The **Transmit Claims** tab is presented.

10. Click Transmit. The **Transmit Claims** window appears.



Transmit Claims


List of claims to transmit

Selected	Claim No	Claim Type	Practitioner	Provider No	Date	Amount
☒	A0056	Medicare	AL	2406901F	04/04/2014	\$36.30
☒	A0060	Medicare	GP	2146681K	07/05/2014	\$326.92
☒	A0067	Medicare	NP1	2406901F	23/11/2016	\$793.73
☒	A0068	Medicare	GP	2406901F	23/05/2014	\$69.00
☒	A0069	Medicare	56	2406901F	05/06/2014	\$70.30
☒	A0070	Medicare	GP	0724479L	05/06/2014	\$42.30
☒	A0073	Medicare	GP	2525213W	24/07/2014	\$92.10
☒	A0074	Medicare	NP1	2406901F	25/07/2014	\$37.05
☒	A0076	Medicare	NP1	2406901F	05/08/2014	\$70.30
☒	A0077	Medicare	AL	2406901F	11/08/2014	\$263.60

Select All Clear All Transmit Cancel

11. Locate and tick the item(s) for transmitting.

12. Click **Transmit**.



1 Select Claims 2 Prepare Claim Data 3 Print Claim Reports 4 Transmit Claims 5 Request Reports

Processing:

*** START of TRANSMISSION at 13/08/2013 9:44:55 AM
 Claim A000001 successfully transmitted at 13/08/2013 9:44:56 AM
 *** END of TRANSMISSION - 1 claim(s) out of 1 sent to Medicare Australia.

Transmit Next Step

13. Click **OK**.

APPENDIX E- RETIRED AND SUSPENDED INDICATORS

PI04 – Fully Immunised Children (SUSPENDED)

Description

Proportion of Indigenous children who are regular patients, aged:

- 12 months to less than 24 months.
- 24 months to less than 36 months.
- 60 months to less than 72 months.

and who are 'fully immunised'.

Business Rules:

For a 12-month-old child to be considered 'fully Immunised' their record must show the following:

	Minimum Count	Antigens
		Each of the antigens have been immunised for times equal or larger than the minimum count. Not to sum up all in the same group
DTPa count	3	DIPHTHERIA, TETANUS, PERTUSSIS
Polio count	3	POLIO
HiB count	2	HIB
Hep B count	2	HEPB
MMR count	0	MEASLES, MUMPS, RUBELLA

For a 24-month-old child to be considered 'fully Immunised' their record must show the following:

	Minimum Count	Antigens
		Each of the antigens have been immunised for times equal or larger than the minimum count. Not to sum up all in the same group.
DTPa count	3	DIPHTHERIA, TETANUS, PERTUSSIS

Polio count	3	POLIO
HiB count	3	HIB
Hep B count	3	HEPB
MMR count	1	MEASLES, MUMPS, RUBELLA

For a 60-month-old child to be considered 'fully Immunised' their record must show the following:

	Minimum Count	Antigens
		Each of the antigens have been immunised for times equal or larger than the minimum count. Not to sum up all in the same group.
DTPa count	4	DIPHTHERIA, TETANUS, PERTUSSIS
Polio count	4	POLIO
MMR count	2	MEASLES, MUMPS, RUBELLA
HiB count	0	HIB
Hep B count	0	HEPB

Calculation 1: 12 Months to Less Than 24 Months

Numerator

How many regular Indigenous patients were fully immunised and aged from 12 months to less than 24 months.

Denominator

How many of your regular Indigenous patients were aged 12 months to less than 24 months.

Calculation 2: 24 Months to Less Than 36 Months

Numerator

How many regular Indigenous patients were fully immunised and aged from 24 months to less than 36 months.

Denominator

How many of your regular Indigenous patients were aged 24 months to less than 36 months.

Calculation 3: 60 Months to Less Than 72 Months

Numerator

How many patients were fully immunised and aged 60 months to less than 72 months.

Denominator

How many of your regular Indigenous patients were aged 60 months to less than 72 months.

Notes


A child is considered 'fully immunised' if they have received the relevant vaccinations as per the National Immunisation Program Schedule by the specified age milestones. Currently these vaccinations include:

- DTPa (diphtheria, tetanus, pertussis).
- Polio.
- HiB (haemophilus influenzae type B).
- Hep B (hepatitis B).
- MMR (measles, mumps, rubella).

Example – To View or Edit Child Immunisation Records


For correct data entry you will need to follow the below workflow.

1. Open the record of a child patient.
2. Select the 'Imm.' Tab.
3. Right click within the white space to add new Immunisations



The screenshot displays the MedicalDirector Clinical 4.2 interface. The patient record is for a 15-month-old male child, Aa. Bankstown, New 2200, with a DOB of 01/01/2023. The 'Imm.' tab is selected in the top navigation bar. A right-click context menu is open over the immunisation table, showing options: 'Record Encounter', 'Delete item', 'Print', 'Refresh', 'Send To AIR', and 'Edit item'. The table has columns: Date, Immunisation, Sequence, Batch, Vaccine serial number, Consent, Provided by, Vaccinator, Site, Comment, and Send to AIR Status. On the left, there are sections for 'AIR Web Services' (including a verification check and notifications) and 'Record Medical Exemption' and 'Record Planned Catch Up' buttons.

4. Add immunisation details and click **Save**



Vaccination Window

☐ Vaccine given elsewhere
 ☐ Vaccine administered overseas
 ☒ Send to AIR

Vaccinator: Dr A Practitioner (MedicalDirector Samples Database)

Submit on behalf of: Dr A Practitioner

Consent provided by: Patient
☐ Vaccination declined

Date: 30/04/2024

Type:

Site:

Sequence:

Batch No: ☐ Store batch No

Vaccine serial no:

Comment: Scan barcode

Mark for recall Save Cancel

Check

For each age group, you cannot have more immunised children than the total number of Indigenous children who were your regular clients.

Note: This indicator is currently suspended - it will not be visible to health services in the health data portal from June 2021 onwards

PI08 – Team Care Arrangement (MBS Item 723) (RETIRED)

Description

Proportion of regular patients who are Indigenous, have a chronic disease (type 2 diabetes) and for whom a Team Care Arrangement (MBS Item 723) was claimed within the previous 24 months.

Currently, Type 2 diabetes is the only chronic disease included for this indicator.

Business Rules

- Patient has a history of Type 2 Diabetes as defined in PI05.
- Patient has had an MBS Item 723 claimed as defined in PI03.

Numerator

The number of regular Indigenous patients that had a Team Care Arrangement (claimed under MBS Item 723) in the **24 months up to the census date**.

In the Numerator, only those patients whose Team Care Arrangement was claimed by your organisation under MBS Item 723.

Denominator

The number of regular Indigenous patients have type 2 diabetes.

Reporting by Age Group

	Age (years)							
	0 to <5	5 to <15	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓	✓	✓

Records Excluded

Patients with type 1 diabetes, secondary diabetes, gestational diabetes mellitus (GDM), previous GDM, impaired fasting glucose or impaired glucose tolerance.

Notes

- The extract only looks for the MBS claim and whether the patient is in the specified disease category (as per the denominator).
- If your organisation does not claim this MBS item but provides an equivalent level of care, such as an equivalent team care approach that cannot be claimed through the MBS.
- A note in the submission comments if your patients cannot access Team Care Arrangements. This is often the case in remote regions.
- Services taking part in the Health Care Homes Trial: For the duration of the Health Care Homes trial (currently 1 October 2017 to 30 November 2019), patients who are part of the trial will be deemed to have had an MBS Item 723 claimed if there is evidence of a Team Care Arrangement recorded.
- While Type 2 Diabetes Is the only chronic disease needed for this nKPI, MedicalDirector Insights also reports on: Cardiovascular disease, Chronic Obstructive Pulmonary Disease and Chronic Kidney Disease.

Example – Creating and Processing a 723 Claim

The method for creating and processing a 723 claim can be seen in PI03.

Example – Recording History of Type 2 Diabetes

A history of Type 2 Diabetes may be added to the patient file by following the method outlined in PI05

Check

For each age and sex at birth group:

- You cannot have more patients for whom an MBS Item 723 (for managing type 2 diabetes under a Team Care Arrangement) was claimed than there are patients with type 2 diabetes.
- The total number of regular patients with type 2 diabetes should be the same as the total number of regular patients with type 2 diabetes in each relevant age and sex at birth group in these nKPIs: PI05, PI07, PI15, PI18 and PI23.
- You cannot have more people with type 2 diabetes than there are in the same age and sex at birth groups in these nKPIs: PI03 (adults), PI09 and PI16.

Note: This indicator is retired - it is not visible to health services in the health data portal from December 2020 onwards

PI15 – Influenza Immunisation (Type 2 Diabetes or COPD Patients) (RETIRED)

Description

Proportion of regular patients who are Indigenous, aged 15–49, are recorded as having type 2 diabetes and are immunised against influenza, **AND SIMILARLY** for chronic obstructive pulmonary disease (COPD).

Business Rules

- Patients with a history of Type II diabetes defined as per PI05.
- Patient's influenza immunisation history defined as per PI14.
- Patients with a history of COPD are defined as having a condition listed under one of the below DOCLE codes:

bronchit@chro	Bronchitis Chronic
coad	CAL (Chronic Airways Limitation)
coad	Chronic Airways Limitation
coad	Chronic Bronchitis
coad	Chronic Obstructive Airways Disease
coad	Chronic Obstructive Pulmonary Disease
coad	COAD
coad	COAD (Chronic Obstructive Airways Disease)
coad	COPD (Chronic Obstructive Pulmonary Disease)
emph	Emphysema

Calculation 1: Type 2 Diabetes

Numerator

The number of regular Indigenous patients that have type 2 diabetes and had an influenza vaccination in the 12 months up to the census date.

Note: These need to be recorded in the immunisations tab like PI04.

Denominator

The number of regular Indigenous patients who have type 2 diabetes.

Calculation 2: COPD

Numerator

The number of regular Indigenous patients that have COPD and had an influenza vaccination in the 12 months up to the census date.

Note: These need to be recorded in the immunisations tab like PI04.

Denominator

The number of regular Indigenous patients that have COPD.

Reported by Age Group

	Age (years)			
	15 to <25	25 to <35	35 to <45	45 to <50
Male	✓	✓	✓	✓
Female	✓	✓	✓	✓

Records Excluded

Patients in the Numerator if they have not been vaccinated, regardless of the reason (you will still include them in the Denominator if they are aged 15 to <50).

Example - Record Influenza Immunisation

Influenza Immunisation may be recorded through the method explained in PI14.

Example - Record history of Type 2 Diabetes or COPD

A history of Type 2 Diabetes may be added following the method explained in PI05.

Check

For each age and sex at birth group:

- You cannot have more people being immunised than the total number of regular patients with either type 2 diabetes or COPD.
- You cannot have more regular patients with type 2 diabetes or COPD than the total number of regular patients in these nKPIs: PI03 (adults), PI09 and PI16.
- The total number of regular patients with type 2 diabetes should be the same as the total number of regular patients with type 2 diabetes in each relevant age and sex at birth group for these nKPIs: PI05, PI07, PI08, PI18 and PI23 (other than for the 45–49 age group, see below).

For age 45–49:

- The total number of regular patients with type 2 diabetes should be less than or the same as the total number of regular patients with type 2 diabetes in the 45–54 age group for these nKPIs: PI05, PI07, PI08, PI18 and PI23.

Note: This indicator is retired - it is not visible to health services in the health data portal from December 2021 onwards - no change required to CIS reports.

PI17 – AUDIT-C Result Recorded (RETIRED)

Description

Proportion of regular patients who are Indigenous, aged 15 and over and who have had an AUDIT-C result recorded within the previous 24 months with result (score) as one of the following:

- Greater than or equal to 4 in males and 3 in females: OR
- Less than 4 in males and 3 in females.

Business Rules

- AUDIT-C result recorded as defined in [PI16](#).
- The score is defined as the result recorded for the AUDIT-C assessment.

Calculation 1: Males with A Score Equal to Or Greater Than 4

Numerator

The number of regular Indigenous male patients had an AUDIT-C score of greater than or equal to 4 in the 24 months up to the census date.

For correct data entry please see PI16.

Denominator

The number of regular Indigenous male patients that had an AUDIT-C result recorded in the same 24 months.

An AUDIT-C with a result within the specified levels recorded in the last 24 months via the Audit-C Assessment only.

Calculation 2: Males with A Score Less Than 4

Numerator

The number of regular Indigenous male patients that had an AUDIT-C score of less than 4 in the 24 months up to the census date.

For correct data entry please see PI16.

Denominator

The number of males who had an AUDIT-C result recorded in the same 24 months.

An AUDIT-C with a result within the specified levels recorded in last 24 months via the Audit-C Assessment only.

Calculation 3: Females with A Score Equal to Or Greater Than 3

Numerator

The number of regular Indigenous female patients that had an AUDIT-C score of greater than or equal to 3 in the 24 months up to the census date.

For correct data entry please see PI16.

Denominator

The number of regular Indigenous female patients that had an AUDIT-C result recorded in the same 24 months.

An AUDIT-C with a result within the specified levels recorded in the last 24 months via the Audit-C Assessment only.

Calculation 4: Females with A Score Less Than 3

Numerator

The number of regular Indigenous female patients that had an AUDIT-C score of less than 3 in the 24 months up to the census date.

For correct data entry please see PI16.

Denominator

The number of females who had an AUDIT-C result recorded in the same 24 months.

An AUDIT-C with a result within the specified levels recorded in the last 24 months via the Audit-C Assessment only.

Reporting by Age Group

	Age (years)					
	15 to <25	25 to <35	35 to <45	45 to <55	55 to <65	65+
Male	✓	✓	✓	✓	✓	✓
Female	✓	✓	✓	✓	✓	✓

Records Excluded

- Results from any other alcohol use screening tool.


Notes

Where an Indigenous regular patient's AUDIT-C score does not have an assessment date assigned within the Patient Information Record System (PIRS), the AUDIT-C score as recorded in the PIRS should be treated as current (that is, as having been updated within the previous 24 months).

Example - Recording Alcohol Consumption Status

A Patient's alcohol consumption status should be entered following the method outlined in PI16.

If this patient has had an Audit-C assessment recorded in their file before, click 'New Assessment' to add a new record.



Patient Details

Pt. Details Allergies/Adverse Reactions/Warnings Family/Social Hx Notes Smoking **Alcohol** Personal Details

Date of assessment: 10/11/2016

Audit-C Assessment

1. How often do you have a drink containing alcohol?

☐ Never ☒ Monthly or less ☐ 2-4 times a month

☐ 2-3 times a week ☐ 4 or more times a week

2. How many standard drinks containing alcohol do you have on a typical day?

☐ 1 or 2 ☐ 3 or 4 ☐ 5 or 6

☐ 7 to 9 ☒ 10 or more

3. How often do you have six or more drinks on one occasion?

☐ Never ☐ Less than monthly ☐ Monthly

☐ Weekly ☒ Daily or almost daily

Audit-C Total Score: 9

In men a score of 4 or more and in women a score of 3 or more is considered positive, optimal for identifying hazardous drinking or active alcohol use disorders. The guidelines to reduce health risks from drinking alcohol provide further assessment and treatment options.

Patient concerned about drinking?

☐ Yes ☐ No ☒ Don't know

[View Alcohol Guidelines](#) [Reference](#) [New Assessment](#)

Currently displaying data from assessment performed on 10/11/2016. Click 'New Assessment' to conduct a new assessment.

☐ Update address for all family members
☐ Auto-capitalise names

[Save](#) [Cancel](#)

Date	Time	Score	Concerns	Comments
10/11/2016	10:55:22	9	Don't know	No
10/11/2016	09:53:28	2	Don't know	No

Delete

Comments

Check

For each age and sex at birth group:

- You cannot have more regular patients with AUDIT-C results in each range than the total number of regular patients who had an AUDIT-C result recorded.
- The total number of regular patients who had an AUDIT-C result recorded should be less than or equal to the number of regular patients who had alcohol consumption status recorded in PI16.

Note: This indicator is retired - it is not visible to health services in the health data portal from December 2022 onwards - no change required to CIS reports.